arXiv:0711.3968v1 [math.GT] 26 Nov 2007

The classification and the conjugacy classes of the finite subgroups of the sphere braid groups

DACIBERG LIMA GONÇALVES
Departamento de Matemática - IME-USP,
Caixa Postal 66281 - Ag. Cidade de São Paulo,
CEP: 05311-970 - São Paulo - SP - Brazil.
e-mail: `dlgoncal@ime.usp.br`

JOHN GUASCHI
Laboratoire de Mathématiques Nicolas Oresme UMR CNRS 6139,
Université de Caen BP 5186,
14032 Caen Cedex, France.
e-mail: `guaschi@math.unicaen.fr`

21st November 2007

Abstract

Let $n \geqslant 3$. We classify the finite groups which are realised as subgroups of the sphere braid group $B_n(\mathbb{S}^2)$. Such groups must be of cohomological period 2 or 4. Depending on the value of n, we show that the following are the maximal finite subgroups of $B_n(\mathbb{S}^2)$: $\mathbb{Z}_{2(n-1)}$; the dicyclic groups of order $4n$ and $4(n-2)$; the binary tetrahedral group T_1; the binary octahedral group O_1; and the binary icosahedral group I. We give geometric as well as some explicit algebraic constructions of these groups in $B_n(\mathbb{S}^2)$, and determine the number of conjugacy classes of such finite subgroups. We also reprove Murasugi's classification of the torsion elements of $B_n(\mathbb{S}^2)$, and explain how the finite subgroups of $B_n(\mathbb{S}^2)$ are related to this classification, as well as to the lower central and derived series of $B_n(\mathbb{S}^2)$.

1 Introduction

The braid groups B_n of the plane were introduced by E. Artin in 1925 [A1, A2]. Braid groups of surfaces were studied by Zariski [Z]. They were later generalised by Fox to braid groups of arbitrary topological spaces via the following definition [FoN]. Let M be a compact, connected surface, and let $n \in \mathbb{N}$. We denote the set of all ordered n-tuples of distinct points of M, known as the *n configuration space of M*, by:

$$F_n(M) = \left\{ (p_1, \ldots, p_n) \;\middle|\; p_i \in M \text{ and } p_i \neq p_j \text{ if } i \neq j \right\}.$$

2000 AMS Subject Classification: 20F36 (primary), 20F50, 20E45, 57M99 (secondary).

Configuration spaces play an important rôle in several branches of mathematics and have been extensively studied, see [CG, FH] for example.

The symmetric group S_n on n letters acts freely on $F_n(M)$ by permuting coordinates. The corresponding quotient will be denoted by $D_n(M)$. The *n pure braid group $P_n(M)$* (respectively the *n braid group $B_n(M)$*) is defined to be the fundamental group of $F_n(M)$ (respectively of $D_n(M)$).

Together with the real projective plane $\mathbb{R}P^2$, the braid groups of the 2-sphere $\mathbb{S}^2$ are of particular interest, notably because they have non-trivial centre [GVB, GG1], and torsion elements [VB, Mu]. Indeed, Van Buskirk showed that among the braid groups of compact, connected surfaces, $B_n(\mathbb{S}^2)$ and $B_n(\mathbb{R}P^2)$ are the only ones to have torsion [VB]. Let us recall briefly some of the properties of $B_n(\mathbb{S}^2)$ [FVB, GVB, VB].

If $\mathbb{D}^2 \subseteq \mathbb{S}^2$ is a topological disc, there is a group homomorphism $\iota\colon B_n \longrightarrow B_n(\mathbb{S}^2)$ induced by the inclusion. If $\beta \in B_n$, we shall denote its image $\iota(\beta)$ simply by β. Then $B_n(\mathbb{S}^2)$ is generated by $\sigma_1, \ldots, \sigma_{n-1}$ which are subject to the following relations:

$$\begin{gathered}
\sigma_i\sigma_j = \sigma_j\sigma_i \text{ if } |i-j| \geqslant 2 \text{ and } 1 \leqslant i,j \leqslant n-1\\
\sigma_i\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\sigma_{i+1} \text{ for all } 1 \leqslant i \leqslant n-2, \text{ and}\\
\sigma_1\cdots\sigma_{n-2}\sigma_{n-1}^2\sigma_{n-2}\cdots\sigma_1 = 1.
\end{gathered}$$

Consequently, $B_n(\mathbb{S}^2)$ is a quotient of B_n. The first three sphere braid groups are finite: $B_1(\mathbb{S}^2)$ is trivial, $B_2(\mathbb{S}^2)$ is cyclic of order 2, and $B_3(\mathbb{S}^2)$ is a ZS-metacyclic group (a group whose Sylow subgroups, commutator subgroup and commutator quotient group are all cyclic) of order 12, isomorphic to the semi-direct product $\mathbb{Z}_3 \rtimes \mathbb{Z}_4$ of cyclic groups, the action being the non-trivial one, which in turn is isomorphic to the dicyclic group Dic_{12} of order 12. The Abelianisation of $B_n(\mathbb{S}^2)$ is isomorphic to the cyclic group $\mathbb{Z}_{2(n-1)}$. The kernel of the associated projection $\xi\colon B_n(\mathbb{S}^2) \longrightarrow \mathbb{Z}_{2(n-1)}$ (which is defined by $\xi(\sigma_i) = \overline{1}$ for all $1 \leqslant i \leqslant n-1$) is the commutator subgroup $\Gamma_2\left(B_n(\mathbb{S}^2)\right)$. If $w \in B_n(\mathbb{S}^2)$ then $\xi(w)$ is the exponent sum (relative to the σ_i) of w modulo $2(n-1)$.

Gillette and Van Buskirk showed that if $n \geqslant 3$ and $k \in \mathbb{N}$ then $B_n(\mathbb{S}^2)$ has an element of order k if and only if k divides one of $2n$, $2(n-1)$ or $2(n-2)$ [GVB]. The torsion elements of $B_n(\mathbb{S}^2)$ and $B_n(\mathbb{R}P^2)$ were later characterised by Murasugi [Mu]. For $B_n(\mathbb{S}^2)$, these elements are as follows:

THEOREM 1 ([Mu]). *Let $n \geqslant 3$. Then the torsion elements of $B_n(\mathbb{S}^2)$ are precisely powers of conjugates of the following three elements:*

(a) $\alpha_0 = \sigma_1\cdots\sigma_{n-2}\sigma_{n-1}$ (which is of order $2n$).

(b) $\alpha_1 = \sigma_1\cdots\sigma_{n-2}\sigma_{n-1}^2$ (of order $2(n-1)$).

(c) $\alpha_2 = \sigma_1\cdots\sigma_{n-3}\sigma_{n-2}^2$ (of order $2(n-2)$).

The three elements α_0, α_1 and α_2 are respectively n, $(n-1)$ and $(n-2)$ roots of Δ_n, where Δ_n is the so-called 'full twist' braid of $B_n(\mathbb{S}^2)$, defined by $\Delta_n = (\sigma_1\cdots\sigma_{n-1})^n$. So $B_n(\mathbb{S}^2)$ admits finite cyclic subgroups isomorphic to $\mathbb{Z}_{2n}$, $\mathbb{Z}_{2(n-1)}$ and $\mathbb{Z}_{2(n-2)}$. In [GG2], we showed that $B_n(\mathbb{S}^2)$ is generated by α_0 and α_1. If $n \geqslant 3$, Δ_n is the unique element of $B_n(\mathbb{S}^2)$ of order 2, and it generates the centre of $B_n(\mathbb{S}^2)$. It is also the square of the *Garside element* (or 'half twist') defined by:

$$T_n = (\sigma_1\cdots\sigma_{n-1})(\sigma_1\cdots\sigma_{n-2})\cdots(\sigma_1\sigma_2)\sigma_1.$$

For $n \geqslant 4$, $B_n(\mathbb{S}^2)$ is infinite. It is an interesting question as to which finite groups are realised as subgroups of $B_n(\mathbb{S}^2)$ (apart of course from the cyclic groups $\langle \alpha_i \rangle$ and their subgroups given in Theorem 1). Another question is the following: how many conjugacy classes are there in $B_n(\mathbb{S}^2)$ of a given abstract finite group? As a partial answer to the first question, we proved in [GG2] that $B_n(\mathbb{S}^2)$ contains an isomorphic copy of the finite group $B_3(\mathbb{S}^2)$ of order 12 if and only if $n \not\equiv 1 \bmod 3$.

While studying the lower central and derived series of the sphere braid groups, we showed that $\Gamma_2\left(B_4(\mathbb{S}^2)\right)$ is isomorphic to a semi-direct product of $\mathcal{Q}_8$ by a free group of rank 2 [GG3]. After having proved this result, we noticed that the question of the realisation of $\mathcal{Q}_8$ as a subgroup of $B_n(\mathbb{S}^2)$ had been explicitly posed by R. Brown [ATD] in connection with the Dirac string trick [F, N] and the fact that the fundamental group of SO(3) is isomorphic to $\mathbb{Z}_2$. The case $n = 4$ was studied by J. G. Thompson [ThJ]. In a previous paper, we provided a complete answer to this question:

THEOREM 2 ([GG4]). *Let $n \in \mathbb{N}$, $n \geqslant 3$.*

(a) $B_n(\mathbb{S}^2)$ contains a subgroup isomorphic to $\mathcal{Q}_8$ if and only if n is even.
(b) If n is divisible by 4 then $\Gamma_2\left(B_n(\mathbb{S}^2)\right)$ contains a subgroup isomorphic to $\mathcal{Q}_8$.

As we also pointed out in [GG4], for all $n \geqslant 3$, the construction of $\mathcal{Q}_8$ may be generalised in order to obtain a subgroup $\langle \alpha_0, T_n \rangle$ of $B_n(\mathbb{S}^2)$ isomorphic to the dicyclic group Dic_{4n} of order $4n$.

It is thus natural to ask which other finite groups are realised as subgroups of $B_n(\mathbb{S}^2)$. One common property of the above subgroups is that they are finite periodic groups of cohomological period 2 or 4. In fact, this is true for all finite subgroups of $B_n(\mathbb{S}^2)$. Indeed, by [GG2], the universal covering X of $F_n(\mathbb{S}^2)$ is a finite-dimensional complex which has the homotopy type of $\mathbb{S}^3$ (we were recently informed by V. Lin that X is biholomorphic to the direct product of $\mathrm{SL}(2,\mathbb{C})$ by the Teichmüller space of the n-punctured Riemann sphere [Li]). Thus any finite subgroup of $B_n(\mathbb{S}^2)$ acts freely on X, and so has period 2 or 4 by Proposition 10.2, Section 10, Chapter VII of [Br]. Since Δ_n is the unique element of order 2 of $B_n(\mathbb{S}^2)$, and it generates the centre $Z(B_n(\mathbb{S}^2))$, the Milnor property must be satisfied for any finite subgroup of $B_n(\mathbb{S}^2)$. Recall also that a finite periodic group G satisfies the p^2-condition (if p is prime and divides the order of G then G has no subgroup isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$), which implies that a Sylow p-subgroup of G is cyclic or generalised quaternion, as well as the $2p$-condition (each subgroup of order $2p$ is cyclic). The classification of finite periodic groups is given by the Suzuki-Zassenhaus theorem (see [AM, ThC] for example), and thus provides a possible line of attack for the subgroup realisation problem. The periods of the different families of these groups were determined in a series of papers by Golasiński and Gonçalves [GoG1, GoG2, GoG3, GoG4, GoG5, GoG6], and so in theory we may obtain a list of those of period 4. A list of all periodic groups of period 4 is provided in [ThC]. However, in the current context, a more direct approach is obtained via the relationship between the braid groups and the mapping class groups of $\mathbb{S}^2$, which we shall now recall.

For $n \in \mathbb{N}$, let $\mathcal{M}_{0,n}$ denote the mapping class group of the n-punctured sphere. We allow the n marked points to be permuted. If $n \geqslant 2$, a presentation of $\mathcal{M}_{0,n}$ is obtained from that of $B_n(\mathbb{S}^2)$ by adding the relation $\Delta_n = 1$ [Ma, MKS]. In other words, we have the following central extension:

$$1 \longrightarrow \langle \Delta_n \rangle \longrightarrow B_n(\mathbb{S}^2) \xrightarrow{p} \mathcal{M}_{0,n} \longrightarrow 1. \tag{1}$$

If $n = 2$, $B_2(\mathbb{S}^2) \cong \mathcal{M}_{0,2} \cong \mathbb{Z}_2$. For $n = 3$, since $\mathcal{M}_{0,3} \cong S_3$, this short exact sequence does not split, and in fact for $n \geqslant 4$ it does not split either [GVB].

This exact sequence may also be obtained in the following manner [Bi]. Let $\mathrm{Diff}^+(\mathbb{S}^2)$ denote the group of orientation-preserving homeomorphisms of $\mathbb{S}^2$, and let $X \in D_n(\mathbb{S}^2)$. Then $\mathrm{Diff}^+(\mathbb{S}^2, X) = \{ f \in \mathrm{Diff}^+(\mathbb{S}^2) \mid f(X) = X \}$ is a subgroup of $\mathrm{Diff}^+(\mathbb{S}^2)$, and we have a fibration $\mathrm{Diff}^+(\mathbb{S}^2, X) \longrightarrow \mathrm{Diff}^+(\mathbb{S}^2) \longrightarrow D_n(\mathbb{S}^2)$, where the basepoint of $D_n(\mathbb{S}^2)$ is taken to be X, and where the second map evaluates an element of $\mathrm{Diff}^+(\mathbb{S}^2)$ on X. The resulting long exact sequence in homotopy yields:

$$\begin{aligned} \cdots \longrightarrow \pi_1\left(\mathrm{Diff}^+\left(\mathbb{S}^2, X\right)\right) \longrightarrow \underbrace{\pi_1\left(\mathrm{Diff}^+\left(\mathbb{S}^2\right)\right)}_{\mathbb{Z}_2} \longrightarrow \underbrace{\pi_1(D_n(\mathbb{S}^2))}_{B_n(\mathbb{S}^2)} \\ \xrightarrow{\delta} \underbrace{\pi_0\left(\mathrm{Diff}^+\left(\mathbb{S}^2, X\right)\right)}_{\mathcal{M}_{0,n}} \longrightarrow \underbrace{\pi_0\left(\mathrm{Diff}^+\left(\mathbb{S}^2\right)\right)}_{=\{1\}}. \end{aligned} \tag{2}$$

The homomorphism $\delta\colon B_n(\mathbb{S}^2) \longrightarrow \mathcal{M}_{0,n}$ is the boundary operator which we shall use in Section 3 in order to describe the geometric realisation of the finite subgroups of $B_n(\mathbb{S}^2)$. If $n \geqslant 3$ then $\pi_1\left(\mathrm{Diff}^+(\mathbb{S}^2, X)\right) = \{1\}$ [EE, Hm, Sc], and we thus recover equation (1) (the interpretation of the Dirac string trick in terms of the sphere braid groups [F, Hn, N] gives rise to the identification of $\pi_1\left(\mathrm{Diff}^+(\mathbb{S}^2)\right)$ with $\langle \Delta_n \rangle$).

In a recent paper, Stukow applies Kerckhoff's solution of the Nielsen realisation problem [K] to classify the finite maximal subgroups of $\mathcal{M}_{0,n}$ [St]. Applying his results to equation (1), we shall see in Section 2 that their counterparts in $B_n(\mathbb{S}^2)$ are cyclic, dicyclic and binary polyhedral groups:

THEOREM 3. *Let $n \geqslant 3$. The maximal finite subgroups of $B_n(\mathbb{S}^2)$ are:*

(a) $\mathbb{Z}_{2(n-1)}$ *if $n \geqslant 5$.*
(b) the dicyclic group Dic_{4n} of order $4n$.
(c) the dicyclic group $\mathrm{Dic}_{4(n-2)}$ if $n = 5$ or $n \geqslant 7$.
(d) the binary tetrahedral group, denoted by T_1, if $n \equiv 4 \bmod 6$.
(e) the binary octahedral group, denoted by O_1, if $n \equiv 0, 2 \bmod 6$.
(f) the binary icosahedral group, denoted by I, if $n \equiv 0, 2, 12, 20 \bmod 30$.

REMARKS 4.

(a) If n is odd then the only finite subgroups of $B_n(\mathbb{S}^2)$ are cyclic or dicyclic. In the latter case, the dicyclic group Dic_{4n} (resp. $\mathrm{Dic}_{4(n-2)}$) is ZS-metacyclic [CM], and is isomorphic to $\mathbb{Z}_n \rtimes \mathbb{Z}_4$ (resp. $\mathbb{Z}_{n-2} \rtimes \mathbb{Z}_4$), where the action is multiplication by -1.
(b) If n is even then one of the binary tetrahedral or octahedral groups is realised as a maximal finite subgroup of $B_n(\mathbb{S}^2)$. Further, since T_1 is a subgroup of O_1, T_1 is realised as a subgroup of $B_n(\mathbb{S}^2)$ for all n even, $n \geqslant 4$.
(c) The groups of Theorem 3 and their subgroups are the finite groups of quaternions [Co]. Indeed, for $p, q, r \in \mathbb{N}$, let us denote

$$\langle p, q, r \rangle = \langle A, B, C \mid A^p = B^q = C^r = ABC \rangle.$$

Then $\mathbb{Z}_{2(n-1)} = \langle n-1, n-1, 1\rangle$, $\mathrm{Dic}_{4n} = \langle n, 2, 2\rangle$, $\mathrm{Dic}_{4(n-2)} = \langle n-2, 2, 2\rangle$, $T_1 = \langle 3, 3, 2\rangle$, $O_1 = \langle 4, 3, 2\rangle$ and $I = \langle 5, 3, 2\rangle$. It is shown in [Co, CM] that for T_1, O_1 and I, this presentation is equivalent to:

$$\langle p, 3, 2\rangle = \left\langle A, B \;\middle|\; A^p = B^3 = (AB)^2 \right\rangle,$$

for $p \in \{3, 4, 5\}$, and that the element A^p is central and is the unique element of order 2 of $\langle p, 3, 2\rangle$.

In Section 2, we also generalise another result of Stukow concerning the conjugacy classes of finite subgroups of $\mathcal{M}_{0,n}$ to $B_n(\mathbb{S}^2)$:

PROPOSITION 5.
(a) Two maximal finite subgroups of $B_n(\mathbb{S}^2)$ are isomorphic if and only if they are conjugate.
(b) Each abstract finite subgroup G of $B_n(\mathbb{S}^2)$ is realised as a single conjugacy class within $B_n(\mathbb{S}^2)$, with the exception, when n is even, of the following cases, for which there are precisely two conjugacy classes:
(i) $G = \mathbb{Z}_4$.
(ii) $G = \mathrm{Dic}_{4r}$, where r divides $\frac{n}{2}$ or $\frac{n-2}{2}$.

In Section 3, we explain how to obtain geometrically the subgroups of Theorem 3, and we also give explicit group presentations of the cyclic and dicyclic subgroups, as well as in the special case T_1 for $n = 4$.

In order to understand better the finite subgroups of $B_n(\mathbb{S}^2)$, it is often useful to know their relationship with the three classes of elements described in Theorem 1. This shall be carried out in Proposition 15 (see Section 4).

The two conjugacy classes of part (b)(i) are realised by the subgroups $\left\langle \alpha_0^{n/2} \right\rangle$ and $\left\langle \alpha_2^{(n-2)/2} \right\rangle$ (they are non conjugate since they project to non-conjugate subgroups in S_n). In Section 5, we construct the two conjugacy classes of part (b)(ii) of Proposition 5:

THEOREM 6. *Let $n \geqslant 4$ be even. Let $N \in \{n, n-2\}$, and let $x = \alpha_0$ (resp. $x = \alpha_0\alpha_2\alpha_0^{-1}$) if $N = n$ (resp. $N = n-2$). Set $N = 2^l k$, where $l \in \mathbb{N}$, and k is odd. Then for $j = 0, 1, \ldots, l$, and q a divisor of k, we have:*

(a) $B_n(\mathbb{S}^2)$ contains 2^j copies of $\mathrm{Dic}_{2^{l+2-j}k/q}$ of the form $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle$, where $i = 0, 1, \ldots, 2^j - 1$.
(b) if $0 \leqslant i, i' \leqslant 2^j - 1$, $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle$ and $\left\langle x^{2^j q}, x^{i'q} T_n \right\rangle$ are conjugate if and only if $i - i'$ is even.

Another question arising from Theorem 2 is the existence of copies of $\mathcal{Q}_8$ lying in $\Gamma_2(B_n(\mathbb{S}^2))$. More generally, one may ask whether the dicyclic groups constructed above (and indeed the other finite subgroups of $B_n(\mathbb{S}^2)$) are contained in $\Gamma_2(B_n(\mathbb{S}^2))$. In the dicyclic case, we have the following result, also proved in Section 5:

PROPOSITION 7. *Let $n \geqslant 4$ be even, let $N \in \{n, n-2\}$, and let r divide N. If r does not divide $N/2$ then the subgroups of $B_n(\mathbb{S}^2)$ abstractly isomorphic to Dic_{4r} are not contained in $\Gamma_2(B_n(\mathbb{S}^2))$. If r divides $N/2$ then up to conjugacy, $B_n(\mathbb{S}^2)$ has a two subgroups abstractly isomorphic to Dic_{4r}, one of which is contained in $\Gamma_2(B_n(\mathbb{S}^2))$, and the other not. In particular, $B_n(\mathbb{S}^2)$ exhibits the two conjugacy classes of $\mathcal{Q}_8$, one of which lies in $\Gamma_2(B_n(\mathbb{S}^2))$, the other not.*

The corresponding result for the binary polyhedral groups may be found in Proposition 16. As a corollary of our results we obtain an alternative proof of Theorem 1 (see Section 6).

Acknowledgements

This work took place during the visits of the second author to the Departmento de Matemática do IME-Universidade de São Paulo during the periods 10th – 27th August 2007 and 4th – 23rd October 2007, and of the visit of the first author to the Laboratoire de Mathématiques Emile Picard, Université Paul Sabatier, Toulouse, during the period 27th August – 17th September 2007. It was supported by the international Cooperation USP/Cofecub project number 105/06, by an 'Aide Ponctuelle de Coopération' from the Université Paul Sabatier, by the Pró-Reitoria de Pesquisa Projeto 1, and by the Projeto Temático FAPESP no. 2004/10229-6 'Topologia Algébrica, Geometrica e Diferencial'.

2 The classification of the finite maximal subgroups of $B_n(\mathbb{S}^2)$

In this section, we prove Theorem 3. We start by making some remarks concerning the central extension (1).

REMARKS 8. Let G be a finite subgroup of $B_n(\mathbb{S}^2)$.
(a) If H is a finite subgroup of $\mathcal{M}_{0,n}$ then $p^{-1}(H)$ is a finite subgroup of $B_n(\mathbb{S}^2)$ of order $2\,|H|$.
(b) If $|G|$ is odd then $\Delta_n \notin G$, and so $G \cong p(G)$. Conversely, if $G \cong p(G)$ then $p|_G$ is injective, and thus $\Delta_n \notin G$, so $|G|$ is odd.
(c) If $|G|$ is even then $\Delta_n \in G$, and so we obtain the following short exact sequence:

$$1 \longrightarrow \langle \Delta_n \rangle \longrightarrow G \xrightarrow{p|_G} p(G) \longrightarrow 1, \tag{3}$$

where $p(G)$ is a finite subgroup of $\mathcal{M}_{0,n}$ of order $\dfrac{|G|}{2}$.
(d) If G is a maximal finite subgroup of $B_n(\mathbb{S}^2)$ then $|G|$ is even, and $p(G)$ is a maximal finite subgroup of $\mathcal{M}_{0,n}$. Conversely, if H is a maximal finite subgroup of $\mathcal{M}_{0,n}$ then $p^{-1}(H)$ is a maximal finite subgroup of $B_n(\mathbb{S}^2)$.

We recall Stukow's theorem:

THEOREM 9 ([St]). *Let $n \geqslant 3$. The maximal finite subgroups of $\mathcal{M}_{0,n}$ are:*
(a) $\mathbb{Z}_{n-1}$ *if* $n \neq 4$.
(b) *the dihedral group* D_{2n} *of order* $2n$.
(c) *the dihedral group* $D_{2(n-2)}$ *if* $n = 5$ *or* $n \geqslant 7$.
(d) A_4 *if* $n \equiv 4, 10 \bmod 12$.
(e) S_4 *if* $n \equiv 0, 2, 6, 8, 12, 14, 18, 20 \bmod 24$.
(f) A_5 *if* $n \equiv 0, 2, 12, 20, 30, 32, 42, 50 \bmod 60$.

REMARK 10. In the case $n = 3$, $\mathcal{M}_{0,3}$ is isomorphic to D_6, obtained as a maximal subgroup in part (b) of Theorem 9, and so its subgroup isomorphic to $\mathbb{Z}_2$ is not maximal. This explains the discrepancy between the value of n in part (a) of Theorems 3 and 9.

Proof of Theorem 3. By Remarks 8, we just need to check that the given groups are those obtained as extensions of $\langle \Delta_n \rangle$ by the groups of Theorem 9. We start by making some preliminary remarks. Let H be one of the finite maximal subgroups of $\mathcal{M}_{0,n}$, and let G be a finite (maximal) subgroup of $B_n(\mathbb{S}^2)$ of order $2\,|H|$ which fits into the following short exact sequence:

$$1 \longrightarrow \langle \Delta_n \rangle \longrightarrow G \xrightarrow{p|_G} H \longrightarrow 1, \tag{4}$$

where $\Delta_n \in G$ belongs to the centre of G, and is the unique element of G of order 2. Then $G = p^{-1}(H)$, and so is unique.

Suppose that $y \in H$ is of order $k \geqslant 2$. Then y has two preimages in G, of the form x and $x\Delta_n$, say, and x is of order k or $2k$. If k is even then by Remarks 8(c), x must be of order $2k$, $x^k = \Delta_n$ and $\Delta_n \in \langle x \rangle$. If k is odd then x is of order k (resp. $2k$) if and only if $x\Delta_n$ is of order $2k$ (resp. k).

A presentation of G may be obtained by applying standard results concerning the presentation of an extension (see Theorem 1, Chapter 13 of [J]). If H is generated by $h_1, \ldots, h_k$ then G is generated by $g_1, \ldots, g_k, \Delta_n$, where $p(g_i) = h_i$ for $i = 1, \ldots, k$. One relation of G is just $\Delta_n^2 = 1$, that of $\mathrm{Ker}\,(p)$. Since $\mathrm{Ker}\,(p) \subseteq Z(G)$, the remaining relations of G are obtained by rewriting the relators of H in terms of the coset representatives, and expressing the corresponding element in the form Δ_n^{ε}, where $\varepsilon \in \{0, 1\}$.

We consider the six cases of Theorem 9 as follows.

(a) $H \cong \mathbb{Z}_{n-1}$: let y be a generator of H, and let $x \in G$ be such that $p(x) = y$. Then $G = \langle \Delta_n, x \rangle$ and $|G| = 2(n-1)$. If n is odd then $\Delta_n \in \langle x \rangle$, $G = \langle x \rangle$, and x is of order $2(n-1)$. If n is even then $G = \langle x\Delta_n \rangle$ (resp. $G = \langle x \rangle$) if x is of order $n-1$ (resp. $2(n-1)$), and $G \cong \mathbb{Z}_{2(n-1)}$ in both cases.

(b) $H \cong \mathrm{D}_{2n}$: let $y, z \in H$ be such that $o(y) = n$, $o(z) = 2$ and $zyz^{-1} = y^{-1}$, and let $x, w \in G$ be such that $p(x) = y$ and $p(w) = z$. So $G = \langle \Delta_n, x, w \rangle$ and $|G| = 4n$. From above, it follows that $w^2 = \Delta_n$, so $G = \langle x, w \rangle$. If n is even then x is of order $2n$ and $x^n = \Delta_n$. The same result may be obtained if n is odd, replacing x by $x\Delta_n$ if necessary. Further, $wxw^{-1}x \in \mathrm{Ker}\,(p)$. If $wxw^{-1}x = \Delta_n$ then $(wx)^2 = 1$. So either $w = x^{-1}$ or $wx = \Delta_n$, and in both cases we conclude that $G = \langle x \rangle$ which contradicts $|G| = 4n$. Hence $wxw^{-1}x = 1$, and since $|G| = 4n$, G is isomorphic to Dic_{4n}.

(c) $H \cong \mathrm{D}_{2(n-2)}$: the previous argument shows that $G \cong \mathrm{Dic}_{4(n-2)}$.

(d) Suppose that H is isomorphic to one of the remaining groups A_4, S_4 or A_5 of Theorem 9. Let $p = 3$ if $H \cong A_4$, $p = 4$ if $H \cong S_4$, and $p = 5$ if $H \cong A_5$. Then H has a presentation given by [Co, CM]:

$$H = \left\langle u, v \;\middle|\; u^2 = v^3 = (uv)^p = 1 \right\rangle.$$

Let $x, w \in G$ be such that $p(x) = u$ and $p(w) = v$. Then $G = \langle x, w, \Delta_n \rangle$. From above, we must have $x^2 = \Delta_n$. Further, replacing w by $w\Delta_n$, we may suppose that $w^3 = \Delta_n$. If $p = 4$ then $(xw)^p = \Delta_n$, while if $p \in \{3, 5\}$, replacing x by $x\Delta_n$ if necessary, we may suppose that $(xw)^p = \Delta_n$. It is shown in [Co, CM] that $x^2 = w^3 = (xw)^p = \Delta_n$ implies that $\Delta_n^2 = 1$, so G admits a presentation given by:

$$G = \left\langle x, w \;\middle|\; x^2 = w^3 = (xw)^p \right\rangle.$$

Thus $G \cong T_1$ if $p = 3$, $G \cong O_1$ if $p = 4$ and $G \cong I$ if $p = 5$. This completes the proof of the theorem.

□

REMARKS 11. Let G_1, G_2 be finite subgroups of $B_n(\mathbb{S}^2)$.
(a) If they are of odd order then by Remarks 8, G_1 and G_2 are isomorphic if and only if $p(G_1)$ and $p(G_2)$ are isomorphic. So suppose that G_1 and G_2 are of even order. If $p(G_1)$ and $p(G_2)$ are isomorphic then it follows from the construction of Theorem 3 that G_1 and G_2 are isomorphic. Conversely, suppose that G_1 and G_2 are isomorphic via an isomorphism $\alpha\colon G_1 \longrightarrow G_2$. Since Δ_n belongs to both, and is the unique element of order 2, we must have $\alpha(\Delta_n) = \Delta_n$, and thus α induces an isomorphism $\widetilde{\alpha}\colon p(G_1) \longrightarrow p(G_2)$ satisfying $\widetilde{\alpha} \circ p = p \circ \alpha$.
(b) If G_1, G_2 are conjugate then clearly so are $p(G_1)$ and $p(G_2)$. Conversely, suppose that $p(G_1), p(G_2)$ are conjugate subgroups of $\mathcal{M}_{0,n}$. Then there exists $g \in \mathcal{M}_{0,n}$ such that $p(G_2) = gp(G_1)g^{-1}$. If G_1 and G_2 are of even order, the fact that equation (1) is a central extension implies that G_1, G_2 are conjugate. If G_1 and G_2 are of odd order, let $L_i = p^{-1}(p(G_i))$ for $i = 1, 2$. Then $[L_i : G_i] = 2$, and it follows from the even order case that L_1 and L_2 are conjugate in $B_n(\mathbb{S}^2)$. But $L_i = G_i \coprod \Delta_n G_i$, and its odd order elements are precisely those of G_i. So the conjugacy between L_1 and L_2 must send G_1 onto G_2.

We are now able to prove Proposition 5.

Proof of Proposition 5. Part (a) follows from Remarks 8 and 11. To prove part (b), let G_1, G_2 be abstractly isomorphic finite subgroups of $B_n(\mathbb{S}^2)$, and for $i = 1, 2$, let $H_i = p(G_i)$. Then $H_1 \cong H_2$: if the G_i are of odd order then $H_i \cong G_i$, so $H_1 \cong H_2$, while if the G_i are of even order, any isomorphism between them must send $\Delta_n \in G_1$ onto $\Delta_n \in G_2$, and so projects to an isomorphism between the H_i. From Remarks 11(b), G_1 and G_2 are conjugate if and only if H_1 and H_2 are, and so the number of conjugacy classes of subgroups of $B_n(\mathbb{S}^2)$ isomorphic to G_1 is the same as the number of conjugacy classes of subgroups of $\mathcal{M}_{0,n}$ isomorphic to H_1. The result follows from the proof of Theorem 3 by remarking that a subgroup of $\mathcal{M}_{0,n}$ isomorphic to $\mathbb{Z}_2$ (resp. D_{2r}) lifts to a subgroup of $B_n(\mathbb{S}^2)$ which is isomorphic to $\mathbb{Z}_4$ (resp. Dic_{4r}). □

3 Realisation of the maximal finite subgroups of $B_n(\mathbb{S}^2)$

In this section, we analyse the geometric and algebraic realisations of the subgroups given in Theorem 3.

3.1 The algebraic realisation of some finite subgroups of $B_n(\mathbb{S}^2)$

The maximal cyclic and dicyclic subgroups of $B_n(\mathbb{S}^2)$ may be realised as follows:
(a) $\mathbb{Z}_{2(n-1)} \cong \langle \alpha_1 \rangle$.
(b) $\mathrm{Dic}_{4n} \cong \langle \alpha_0, T_n \rangle$ [GG4].
(c) The algebraic realisation of $\mathrm{Dic}_{4(n-2)}$ is given by the following proposition:

PROPOSITION 12. *For all $n \geqslant 3$, the subgroup $\left\langle \alpha_0\alpha_2\alpha_0^{-1}, T_n \right\rangle$ of $B_n(\mathbb{S}^2)$ is isomorphic to $\mathrm{Dic}_{4(n-2)}$.*

Proof. Let $x = \alpha_0\alpha_2\alpha_0^{-1}$. We know that x is of order $2(n-2)$, and that $x^{n-1} = \Delta_n = T_n^2$. Further, by standard properties of the corresponding elements in B_n [Bi], $\alpha_0\sigma_i\alpha_0^{-1} = \sigma_{i+1}$ for all $i = 1, \ldots, n-2$, and $T_n\sigma_iT_n^{-1} = \sigma_{n-i}$ for all $i = 1, \ldots, n-1$. Hence $x = \sigma_2 \cdots \sigma_{n-2}\sigma_{n-1}^2$, and

$$T_nxT_n^{-1} = \sigma_{n-2} \cdots \sigma_2\sigma_1^2 = \sigma_{n-1}^{-2}\sigma_{n-2}^{-1} \cdots \sigma_2^{-1} = x^{-1}.$$

Thus $\langle x, T_n\rangle$ is isomorphic to a quotient of $\mathrm{Dic}_{4(n-2)}$. But $T_n \notin \langle x\rangle$, so $\langle x, T_n\rangle$ contains the $2(n-2)+1$ distinct elements of $\langle x\rangle \cup \{T_n\}$, and the result follows. ☐

REMARK 13. In the special case $n = 4$, the binary tetrahedral group T_1 may be realised as follows. Let $y = \sigma_1\sigma_3^{-1}$. From [GG4], we know that $\langle y, T_4\rangle \cong \mathcal{Q}_8$. In $B_4(\mathbb{S}^2)$, we also have $(\sigma_2\sigma_1)^3 = (\sigma_2\sigma_3)^3 = \Delta_4 = T_4^2$. Then $\langle \alpha_1^2\rangle \cong \mathbb{Z}_3$ acts on $\langle y, T_4\rangle$ as follows:

$$\begin{aligned}
\alpha_1^2 \cdot T_4 \cdot \alpha_1^{-2} &= \alpha_1^2(T_4\alpha_1^{-2}T_4^{-1})T_4 \\
&= \alpha_1^2(\sigma_1^{-2}\sigma_2^{-1}\sigma_3^{-1})^2T_4 \text{ (by the action of } T_4) \\
&= \alpha_1^2(\sigma_2\sigma_3)^2T_4 \text{ (using the surface relation of } B_n(\mathbb{S}^2)) \\
&= (\sigma_1\sigma_2\sigma_3^2)^2 \cdot \sigma_3^{-1}\sigma_2^{-1} \cdot (\sigma_2\sigma_3)^3T_4 \\
&= \sigma_1\sigma_2\sigma_3\sigma_1\sigma_2\sigma_1 \cdot \sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1} \cdot \sigma_3\sigma_1\sigma_2\sigma_3\sigma_2^{-1}T_4^3 \text{ (as } T_4^2 = (\sigma_2\sigma_3)^3) \\
&= T_4\sigma_1^{-1}\sigma_2^{-1}\sigma_3\sigma_2\sigma_3\sigma_2^{-1}T_4^3 \text{ (as } \sigma_1 \text{ commutes with } \sigma_3) \\
&= T_4\sigma_1^{-1}\sigma_3T_4^3 \text{ (by the Artin braid relations)} \\
&= T_4y^{-1}T_4^{-1} = y \text{ (by the action of } T_4 \text{ on } y).
\end{aligned}$$

Further,

$$\begin{aligned}
\alpha_1^2 \cdot y \cdot \alpha_1^{-2} &= (\sigma_1^{-1}\sigma_2^{-1})^2 \cdot \sigma_1\sigma_3^{-1} \cdot (\sigma_2\sigma_1)^2 \\
&= (\sigma_1^{-1}\sigma_2^{-1})^2 \cdot \sigma_3^{-1}\sigma_2^{-1} \cdot (\sigma_2\sigma_1)^3 \text{ (as } \sigma_1 \text{ commutes with } \sigma_3) \\
&= \sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1} \cdot \sigma_2^{-1}\sigma_3^{-1}\sigma_2^{-1} \cdot T_4^2 \text{ (as } T_4^2 = (\sigma_2\sigma_1)^3) \\
&= \sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1}\sigma_3^{-1}\sigma_2^{-1}\sigma_3^{-1} \cdot T_4^2 \text{ (by the Artin braid relations)} \\
&= \sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1}\sigma_3^{-1}\sigma_2^{-1}\sigma_1^{-1} \cdot \sigma_1\sigma_3^{-1}T_4^2 \\
&= T_4^{-1}yT_4^2 = T_4y \text{ (since } T_4^2 \text{ is central).}
\end{aligned}$$

Hence $T_1 = \mathcal{Q}_8 \rtimes \mathbb{Z}_3 \cong \langle y, T_4\rangle \rtimes \langle \alpha_1^2\rangle$.

REMARK 14. We also have an algebraic representation of T_1 in $B_6(\mathbb{S}^2)$. Let

$$\begin{aligned}
\gamma &= \sigma_5\sigma_4\sigma_1^{-1}\sigma_2^{-1}, \text{ and} \\
\delta &= \sigma_3^{-1}\sigma_4^{-1}\sigma_5^{-1}\left(\sigma_2^{-1}\sigma_1^{-1}\sigma_2^{-1}\right)\sigma_5\sigma_4\sigma_3.
\end{aligned}$$

Then we claim that $\langle \gamma, \delta\rangle \cong \mathcal{Q}_8 \rtimes \mathbb{Z}_3 \cong T_1$, where the action permutes the elements i, j, k of $\mathcal{Q}_8$. First, $\gamma^3 = \delta^2 = \Delta_6$. We now consider the subgroup $H = \langle \delta, \gamma\delta\gamma^{-1}\rangle$. The action of conjugation by γ permutes cyclically the elements δ, $\gamma\delta\gamma^{-1}$ and $\gamma\delta^2\gamma^{-1}$, so is compatible with the action of $\mathbb{Z}_3$ on $\mathcal{Q}_8$. It just remains to show that $H \cong \mathcal{Q}_8$. Clearly $\delta^2 = (\gamma\delta\gamma^{-1})^2 = \Delta_6$. Let us now prove that

$$\delta^{-1} \cdot \gamma\delta\gamma^{-1} \cdot \delta = \gamma\delta^{-1}\gamma^{-1}. \tag{5}$$

Set $\rho = \sigma_5\sigma_4\sigma_3$, $\gamma' = \rho\gamma\rho^{-1}$ and $\delta' = \rho\delta\rho^{-1}$. Then equation (5) is in turn equivalent to:

$$\begin{aligned}
&\delta'^{-1} \cdot \gamma'\delta'\gamma'^{-1} \cdot \delta' = \gamma'\delta'^{-1}\gamma'^{-1}\\
&\delta'^{-1}\gamma'\delta'\gamma'^{-1}\delta'^{2}\delta'^{-1}\gamma'\delta'\gamma'^{-1} = 1\\
&[\delta'^{-1}, \gamma']^2 = \delta'^{-2} = \Delta_6.
\end{aligned}$$

We shall show that the latter relation holds. Notice that

$$\gamma' = \sigma_5\sigma_4\sigma_3\sigma_5\sigma_4\sigma_1^{-1}\sigma_2^{-1}\sigma_3^{-1}\sigma_4^{-1}\sigma_5^{-1} = \sigma_5\sigma_4\sigma_5\sigma_3\sigma_4\alpha_0.$$

Then

$$\begin{aligned}
[\delta'^{-1}, \gamma'] &= \sigma_5^{-1}\sigma_4^{-1}\sigma_5^{-1}\sigma_2\sigma_1\sigma_2 \cdot \sigma_5\sigma_4\sigma_5\sigma_3\sigma_4\alpha_0 \cdot \sigma_2^{-1}\sigma_1^{-1}\sigma_2^{-1}\sigma_5\sigma_4\sigma_5 \cdot \alpha_0^{-1}\sigma_4^{-1}\sigma_3^{-1}\sigma_5^{-1}\sigma_4^{-1}\sigma_5^{-1}\\
&= \sigma_2\alpha_0\sigma_5^{-1}\alpha_0\sigma_2^{-1}\sigma_1^{-1}\sigma_2^{-1}\sigma_5\sigma_4\sigma_5\sigma_5\sigma_4\sigma_3\sigma_2\sigma_1\sigma_4^{-1}\sigma_3^{-1}\sigma_5^{-1}\sigma_4^{-1}\sigma_5^{-1}\\
&= \sigma_2\alpha_0\sigma_5^{-1}\alpha_0\sigma_2^{-1}\sigma_1^{-1}\sigma_2^{-1}\sigma_5\sigma_3^{-1}\sigma_2^{-1}\sigma_1^{-1}\sigma_4^{-1}\sigma_3^{-1}\sigma_5^{-1}\sigma_4^{-1}\sigma_5^{-1}\\
&= \sigma_2\alpha_0\sigma_5^{-1}\alpha_0\sigma_2^{-1}\sigma_5\sigma_1^{-1}\sigma_2^{-1}\sigma_3^{-1}\sigma_4^{-1}\sigma_5^{-1}\sigma_1\sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1}\sigma_3^{-1}\sigma_4^{-1}\sigma_5^{-1}\\
&= \sigma_2\alpha_0\sigma_5^{-1}\alpha_0\sigma_2^{-1}\sigma_5\alpha_0\sigma_1\sigma_2^{-1}\alpha_0\\
&= \sigma_2\alpha_0\sigma_5^{-1}\alpha_0^{-1}\alpha_0^{2}\sigma_2^{-1}\sigma_5\alpha_0^{-2}\alpha_0^{3}\sigma_1\sigma_2^{-1}\alpha_0^{-3}\alpha_0^{4}\\
&= \sigma_2\tau^{-1}\sigma_4^{-1}\sigma_1\sigma_4\sigma_5^{-1}\alpha_0^{4} = \sigma_2\tau^{-1}\sigma_1\sigma_5^{-1}\alpha_0^{4},
\end{aligned}$$

since conjugation by α_0 permutes cyclically the elements $\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5$ and $\tau = \alpha_0\sigma_5\alpha_0^{-1}$. Thus

$$[\delta'^{-1}, \gamma']^2 = \sigma_2\tau^{-1}\sigma_1\sigma_5^{-1}\alpha_0^{4}\sigma_2\tau^{-1}\sigma_1\sigma_5^{-1}\alpha_0^{-4}\alpha_0^{8} = \sigma_2\tau^{-1}\sigma_1\sigma_5^{-1}\tau\sigma_4^{-1}\sigma_5\sigma_3^{-1}\alpha_0^{8}.$$

Let $\xi = \sigma_2\tau^{-1}\sigma_1\sigma_5^{-1}\tau\sigma_4^{-1}\sigma_5\sigma_3^{-1}$. To prove that $[\delta'^{-1}, \gamma']^2 = \Delta_6 = \alpha_0^{6}$, it suffices to show that $\xi\alpha_0^2 = 1$. Now

$$\begin{aligned}
\xi\alpha_0^2 &= \sigma_2\tau^{-1}\sigma_1\sigma_5^{-1}\tau\sigma_4^{-1}\sigma_5\sigma_3^{-1}\alpha_0^{2} = \sigma_2\alpha_0\sigma_5^{-1}\alpha_0^{-1}\sigma_1\sigma_5^{-1}\alpha_0\sigma_5\alpha_0^{-1}\sigma_4^{-1}\sigma_5\sigma_3^{-1}\alpha_0^{2}\\
&= \sigma_2\alpha_0\sigma_5^{-1}\alpha_0\sigma_5\alpha_0^{-1}\sigma_4^{-1}\sigma_5\sigma_3^{-1}\sigma_4\sigma_2^{-1}\alpha_0 = \sigma_2\alpha_0\sigma_5^{-1}\alpha_0\sigma_5\sigma_3^{-1}\sigma_4\sigma_2^{-1}\sigma_3\sigma_1^{-1}\\
&= \sigma_2\sigma_1\sigma_2\sigma_3\sigma_4\sigma_1\sigma_2\sigma_3\sigma_4\sigma_5^{2}\sigma_4\sigma_3\sigma_3^{-1}\sigma_4^{-1}\sigma_3^{-1}\sigma_4\sigma_2^{-1}\sigma_3\sigma_1^{-1}\\
&= \sigma_1\sigma_2\sigma_1\sigma_3\sigma_4\sigma_1^{-1}\sigma_2^{-1}\sigma_4^{-1}\sigma_3^{-1}\sigma_2^{-1}\sigma_3\sigma_1^{-1} = 1.
\end{aligned}$$

This proves the claim, so $\langle \gamma, \delta \rangle \cong T_1$.

3.2 The geometric realisation of the finite subgroups of $B_n(\mathbb{S}^2)$

The geometric realisation of the finite subgroups may be obtained by letting the corresponding subgroup of $\mathcal{M}_{0,n}$ act on the sphere with the n strings attached in an appropriate manner. For the subgroups Dic_{4n}, $\mathbb{Z}_{2(n-1)}$ and $\mathrm{Dic}_{4(n-2)}$, we attach strings to n symmetrically-distributed points (resp. $n-1$, $n-2$ points) on the equator, and 0 (resp. 1, 2) points at the poles. For T_1, O_1 and I, the n strings are attached symmetrically with respect to the associated regular polyhedron (for the values of n given by Theorem 3) in the following manner.

(d) Let $H = A_4$ be the group of orientation-preserving symmetries of the tetrahedron. Then $n = 6k + 4$, $k \geqslant 0$, and we take k equally-spaced points in the interior of each edge, plus one point at each vertex (or face).

(e) Let $H = S_4$ be the group of orientation-preserving symmetries of the cube (or octahedron).

(i) $n = 12k$, $k \in \mathbb{N}$: take k equally-spaced points in the interior of each edge.

(ii) $n = 12k + 2$, $k \in \mathbb{N}$: take $k - 1$ equally-spaced points in the interior of each edge, plus one point at each vertex and on each face.

(iii) $n = 12k + 6$, $k \geqslant 0$: take k equally-spaced points in the interior of each edge, plus one point on each face.

(iv) $n = 12k + 8$, $k \geqslant 0$: take k equally-spaced points in the interior of each edge, plus one point at each vertex.

(f) Let $H = A_5$ be the group of orientation-preserving symmetries of the icosahedron (or dodecahedron), which has 12 faces, 30 edges and 20 vertices.

(i) $n = 30k$, $k \in \mathbb{N}$: take k equally-spaced points in the interior of each edge.

(ii) $n = 30k + 2$, $k \in \mathbb{N}$: take $k - 1$ equally-spaced points in the interior of each edge, plus one point at each vertex and on each face.

(iii) $n = 30k + 12$, $k \geqslant 0$: take k equally-spaced points in the interior of each edge, plus one point on each face.

(iv) $n = 30k + 20$, $k \geqslant 0$: take k equally-spaced points in the interior of each edge, plus one point at each vertex.

In each case, the action of the given group H of symmetries yields the corresponding maximal finite subgroup of $B_n(\mathbb{S}^2)$. This follows essentially from the definition of the boundary operator $\partial\colon \pi_1(D_n(\mathbb{S}^2)) \longrightarrow \pi_0\left(\text{Diff}^+(\mathbb{S}^2, X)\right)$ in the long exact sequence (2) which we now describe in detail in our setting. As in Section 1, let X be the basepoint in $D_n(\mathbb{S}^2)$, and let $\psi\colon\ \text{Diff}^+(\mathbb{S}^2) \longrightarrow D_n(\mathbb{S}^2)$ denote evaluation on X. So if $g \in \text{Diff}^+(\mathbb{S}^2)$ then $\psi(g) = g(X)$. Let $\text{Id}_{\mathbb{S}^2}$ be the basepoint in $\text{Diff}^+(\mathbb{S}^2)$, so that $\psi(\text{Id}_{\mathbb{S}^2}) = X$. Let $\beta \in B_n(\mathbb{S}^2)$ be a braid, and let $f\colon\ [0,1] \longrightarrow D_n(\mathbb{S}^2)$ be a geometric braid which represents β. So $f(0) = f(1) = X$, and the loop class $\langle f \rangle$ in $B_n(\mathbb{S}^2)$ is equal to β. Then f lifts to $\widetilde{f}\colon\ [0,1] \longrightarrow \text{Diff}^+(\mathbb{S}^2)$ which satisfies $\widetilde{f}(0) = \text{Id}_{\mathbb{S}^2}$ and $\psi \circ \widetilde{f} = f$. Hence $\psi \circ \widetilde{f}(1) = f(1) = X$, and thus $\widetilde{f}(1)$ belongs to the fibre $\text{Diff}^+(\mathbb{S}^2, X)$. Geometrically, $\widetilde{f}$ is an isotopy of $\mathbb{S}^2$ which realises β on the points of X. Neither $\widetilde{f}$ nor the corresponding endpoint $\widetilde{f}(1)$ are unique, however all of the possible $\widetilde{f}(1)$ belong to the same connected component of $\text{Diff}^+(\mathbb{S}^2, X)$, and so determine a unique element, denoted $[\widetilde{f}(1)]$, of $\pi_0\left(\text{Diff}^+(\mathbb{S}^2, X)\right)$, which is the image under ∂ of β. Thus if $\widetilde{f}$ is an isotopy of $\mathbb{S}^2$ which realises β, $\partial(\beta)$ is the mapping class of the homeomorphism $\widetilde{f}(1)$, and corresponds geometrically to just remembering the final homeomorphism (in particular, one forgets the strings of β).

Conversely, if $g \in \text{Diff}^+(\mathbb{S}^2)$ satisfies $g(X) = X$, let $h\colon\ [0,1] \longrightarrow \text{Diff}^+(\mathbb{S}^2)$ be an isotopy from $h(0) = \text{Id}_{\mathbb{S}^2}$ to $h(1) = g$. Then $\psi \circ h$ is a loop in $D_n(\mathbb{S}^2)$ based at X, so describes a geometric braid obtained by attaching strings at the points of X and following the isotopy h. In $\mathbb{S}^2 \times [0,1]$, the strings are given by $\{(\psi \circ h(t), t)\}_{t \in [0,1]} = \{(h(t)(X), t)\}_{t \in [0,1]}$. Thus $\langle \psi \circ h \rangle \in B_n(\mathbb{S}^2)$ is a braid, and by the above construction, $\partial(\langle \psi \circ h \rangle) = [h(1)] = [g]$. In other words, a choice of isotopy h between the identity and $g \in \text{Diff}^+(\mathbb{S}^2, X)$ allows us to lift the mapping class $[g]$ to a preimage $\beta = \langle \psi \circ h \rangle$ under ∂ which is obtained geometrically by attaching strings to X during the isotopy h.

Let $r\colon [0,1] \longrightarrow \mathrm{Diff}^+(\mathbb{S}^2)$ denote rigid rotation through an angle 2π. So $r(0) = r(1) = \mathrm{Id}_{\mathbb{S}^2}$, the loop class $\langle r \rangle$ generates $\pi_1\left(\mathrm{Diff}^+(\mathbb{S}^2)\right) \cong \mathbb{Z}_2$, and thus $\langle \psi \circ r \rangle = \psi_*(\langle r \rangle) = \Delta_n$ since $\psi_*\colon \pi_1\left(\mathrm{Diff}^+(\mathbb{S}^2)\right) \longrightarrow B_n(\mathbb{S}^2)$ is injective. The second preimage of $[g]$ under ∂ is obtained by considering the isotopy $h'\colon [0,1] \longrightarrow \mathrm{Diff}^+(\mathbb{S}^2)$ which is the isotopy h followed by r. The braids $\langle \psi \circ h \rangle$ and $\langle \psi \circ h' \rangle$ differ by $\langle \psi \circ r \rangle = \Delta_n$, and thus define the two preimages of $[g]$ under ∂.

Finally, each finite subgroup H of $\mathcal{M}_{0,n}$ is realised by a finite subgroup of isometries of $\mathbb{S}^2$ (which are the finite subgroups of SO(3)) [K]. Each element of H admits two preimages in $B_n(\mathbb{S}^2)$ which differ by Δ_n. These preimages thus make up the finite subgroup $\partial^{-1}(H)$ of $B_n(\mathbb{S}^2)$ whose order is twice that of H.

4 Position of the finite subgroups of $B_n(\mathbb{S}^2)$ relative to Murasugi's classification

Let $n \geqslant 4$ be even. For $i = 0, 1, 2$, let G_i be the set of torsion elements of $B_n(\mathbb{S}^2)$ whose order divides $2(n-i)$. Equivalently, by Theorem 1, G_i is the set of conjugates of powers of α_i. Notice that G_i is invariant under conjugation, $G_i \cap G_j = \langle \Delta_n \rangle$ for all $0 \leqslant i < j \leqslant 2$, and $G_0 \cup G_1 \cup G_2$ is the set of torsion elements of $B_n(\mathbb{S}^2)$. For many purposes, it is often useful to know where a finite subgroup H of $B_n(\mathbb{S}^2)$ lies relative to the G_i. In this section, we carry out this calculation for all such subgroups.

PROPOSITION 15. *Let H be a finite subgroup of $B_n(\mathbb{S}^2)$ of order greater than or equal to* 3.

(I) Suppose that H is cyclic.

(a) if $|H| = 4$ and n is even then there exists a subgroup H' of $B_n(\mathbb{S}^2)$ isomorphic to $\mathbb{Z}_4$ non-conjugate to H. One of H, H' lies in G_0, while the other lies in G_2.

(b) if either $|H| = 4$ and n is odd, or if $|H| \neq 4$ then $H \subset G_i$, where $|H| \mid 2(n-i)$, and $i \in \{0, 1, 2\}$.

(II) Suppose that H is a subgroup of a maximal non-cyclic subgroup of $B_n(\mathbb{S}^2)$.

(a) If H is a non-cyclic subgroup contained in Dic_{4n} or $Dic_{4(n-2)}$ then it is itself dicyclic, of the form Dic_{4k}, where $k > 1$ divides n or $n-2$ respectively. Further:

(i) if n is odd then $H \subset G_i \cup G_1$, where $i \in \{0, 2\}$ and $|H| \mid 4(n-i)$.

(ii) Suppose that n is even.

(1) if $k \mid n$ (resp. $k \mid n-2$) but $k \nmid \frac{n}{2}$ (resp. $k \nmid \frac{n-2}{2}$) then H lies in $G_0 \cup G_2$ and meets both G_0 and G_2.

(2) if $k \mid \frac{n}{2}$ (resp. $k \mid \frac{n-2}{2}$) then there exists another subgroup H' of $B_n(\mathbb{S}^2)$ isomorphic to Dic_{4k} but non conjugate to H. In this case, one of H, H' is contained wholly within G_0 (resp. G_2), and the other lies in $G_0 \cup G_2$ and meets both G_0 and G_2.

(b) Suppose that H is a subgroup of a copy of T_1 in the case that T_1 is maximal.

(i) If $H \cong T_1$ then H lies in $G_0 \cup G_1$ (resp. $G_2 \cup G_1$) if $n \equiv 4 \bmod 12$ (resp. $n \equiv 10 \bmod 12$), and meets both G_0 (resp. G_2) and G_1.

(ii) If H is isomorphic to $\mathbb{Z}_3$ or $\mathbb{Z}_6$ then it is contained in G_1.

(iii) If H is isomorphic to $\mathbb{Z}_4$ or $\mathcal{Q}_8$ then it is contained in G_0 if $n \equiv 4 \bmod 12$, and in G_2 if $n \equiv 10 \bmod 12$.

(c) Suppose that H is a subgroup of a copy of I in the case that I is maximal.

(i) If H is isomorphic to I then H is contained in G_0 (resp. G_2) if $n \equiv 0 \bmod 60$ *(resp.* $n \equiv 2 \bmod 60$*), and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 12, 20, 30, 32, 42, 50 \bmod 60$.
(ii) If H is isomorphic to $\mathbb{Z}_3$ or $\mathbb{Z}_6$ then it is contained in G_0 if $n \equiv 0, 12 \bmod 30$*, and in G_2 if* $n \equiv 2, 20 \bmod 30$.
(iii) If H is isomorphic to $\mathbb{Z}_5$ or $\mathbb{Z}_{10}$ then it is contained in G_0 if $n \equiv 0, 20 \bmod 30$*, and in G_2 if* $n \equiv 2, 12 \bmod 30$.
(iv) If H is isomorphic to $\mathbb{Z}_4$ or $\mathcal{Q}_8$ then it is contained in G_0 if $n \equiv 0, 12, 20, 32 \bmod 60$*, and in G_2 if* $n \equiv 2, 30, 42, 50 \bmod 60$.
(v) If H is isomorphic to T_1 or to Dic_{12} then it lies in G_0 if $n \equiv 0, 12 \bmod 60$*, in G_2 if* $n \equiv 2, 50 \bmod 60$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 20, 30, 32, 42 \bmod 60$.
(vi) If H is isomorphic to Dic_{20} then it lies in G_0 if $n \equiv 0, 20 \bmod 60$*, in G_2 if* $n \equiv 2, 42 \bmod 60$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 12, 30, 32, 50 \bmod 60$.

(d) Suppose that H is a subgroup of a copy of O_1 in the case that O_1 is maximal.
(i) If H is isomorphic to O_1 then it lies in G_0 if $n \equiv 0 \bmod 24$*, in G_2 if* $n \equiv 2 \bmod 24$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 6, 8, 12, 14, 18, 20 \bmod 24$.
(ii) If H is isomorphic to T_1 then it lies in G_0 if $n \equiv 0 \bmod 12$*, in G_2 if* $n \equiv 2 \bmod 12$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 6, 8 \bmod 12$.
(iii) If H is isomorphic to $\mathcal{Q}_{16}$ then it lies in G_0 if $n \equiv 0, 8 \bmod 24$*, in G_2 if* $n \equiv 2, 18 \bmod 24$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 6, 12, 14, 20 \bmod 24$.
(iv) If H is isomorphic to Dic_{12} then it lies in G_0 if $n \equiv 0, 6 \bmod 24$*, in G_2 if* $n \equiv 2, 20 \bmod 24$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 8, 12, 14, 18 \bmod 24$.
(v) If H is isomorphic to $\mathbb{Z}_8$ then it lies in G_0 if $n \equiv 0, 8 \bmod 12$*, and in G_2 if* $n \equiv 2, 6 \bmod 12$.
(vi) If H is isomorphic to $\mathbb{Z}_4$ then there exists another non-conjugate subgroup H' of $B_n(\mathbb{S}^2)$ isomorphic to $\mathbb{Z}_4$. One of H, H' is contained in G_0 if $n \equiv 0, 8 \bmod 12$*, and in G_2 if* $n \equiv 2, 6 \bmod 12$*, while the other is contained in G_0 if* $n \equiv 0, 6, 8, 14 \bmod 24$*, and to G_2 if* $n \equiv 2, 12, 18, 20 \bmod 24$.
(vii) If H is isomorphic to $\mathcal{Q}_8$ then there exists another non-conjugate subgroup H' of $B_n(\mathbb{S}^2)$ isomorphic to $\mathcal{Q}_8$. One of H, H' is contained in G_0 if $n \equiv 0, 8 \bmod 12$*, and to G_2 if* $n \equiv 2, 6 \bmod 12$*, while the other lies in G_0 if* $n \equiv 0, 8 \bmod 24$*, in G_2 if* $n \equiv 2, 18 \bmod 24$*, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if* $n \equiv 6, 12, 14, 20 \bmod 24$.
(viii) If H is isomorphic to $\mathbb{Z}_3$ or $\mathbb{Z}_6$ then it lies in G_0 if $n \equiv 0 \bmod 6$ *and in G_2 if* $n \equiv 2 \bmod 6$.

Proof. Let H be a finite subgroup of $B_n(\mathbb{S}^2)$ of order at least three.
(I) Suppose first that H is cyclic. Since $G_i \cap G_j = \langle \Delta_n \rangle$ and $|\langle \alpha_i \rangle| = 2(n-i)$, the order of H is sufficient to decide where H lies, unless n is even and H is of order 4, in which case there is another non-conjugate subgroup H' isomorphic to $\mathbb{Z}_4$. One of H, H' is conjugate to $\left\langle \alpha_0^{n/2} \right\rangle$ which is contained in G_0, while the other is conjugate to $\left\langle \alpha_2^{(n-2)/2} \right\rangle$ which lies in G_2. These two cases may be distinguished easily by checking the permutation of a generator of H, H'.
(II) Now suppose that H is a subgroup of a maximal non-cyclic subgroup of $B_n(\mathbb{S}^2)$. We consider the possible cases in turn.

(a) Firstly, let H be a subgroup of the dicyclic group Dic_{4n}, which up to conjugation may be assumed to be $\langle \alpha_0, T_n \rangle = \langle \alpha_0 \rangle \coprod T_n \langle \alpha_0 \rangle$. We first suppose that n is odd. Then $\langle \alpha_0 \rangle \subset G_0$, and the coset $T_n \langle \alpha_0 \rangle$ consists of the elements of Dic_{4n} of order 4, so lies in G_1. The group Dic_{4n} fits into a short exact sequence:

$$1 \longrightarrow \mathbb{Z}_n \longrightarrow \mathrm{Dic}_{4n} \xrightarrow{g} \mathbb{Z}_4 \longrightarrow 1.$$

If $g(H) = \{\overline{0}\}$, then $H < \mathbb{Z}_n$, and H is cyclic, of order dividing n, so lies in G_0. If $g(H) = \{\overline{0}, \overline{2}\}$, then $H < \mathbb{Z}_{2n}$, and again H is cyclic, of order dividing $2n$, so lies in G_0. Finally, if $g(H) = \mathbb{Z}_4$ then we have

$$1 \longrightarrow H \cap \mathbb{Z}_n \longrightarrow H \xrightarrow{g} \mathbb{Z}_4 \longrightarrow 1,$$

and $H \cong \mathbb{Z}_k \rtimes \mathbb{Z}_4$, where k divides n. If $k = 1$ then $H \cong \mathbb{Z}_4$. Since n is odd, H must then lie in G_1. So suppose that $k > 1$. Then $H = \left\langle \alpha_0^{n/k}, T_n \right\rangle$ is dicyclic, and so lies in $G_0 \cup G_1$.

Now suppose that n is even. Then Dic_{4n} fits into the following short exact sequence:

$$1 \longrightarrow \mathbb{Z}_{2n} \longrightarrow \mathrm{Dic}_{4n} \xrightarrow{f} \mathbb{Z}_2 \longrightarrow 1.$$

If $f(H) = \{\overline{0}\}$ then $H \subset \mathbb{Z}_{2n}$ and so lies in G_0. If $f(H) = \mathbb{Z}_2$ and $H \cap \mathbb{Z}_{2n}$ were of odd order, then H would be both dicyclic and of order twice an odd number, which cannot occur. So suppose that $f(H) = \mathbb{Z}_2$ and $H \cap \mathbb{Z}_{2n}$ is of even order, $2k$, say, where $k \mid n$. If $k = 1$ then $H \cong \mathbb{Z}_4$, and H may lie in G_0 or G_2 depending on the permutation of its generators. So suppose that $k \geqslant 2$. Then H is dicyclic of order $4k$. Now

$$\mathrm{Dic}_{4n} = \underbrace{\langle \alpha_0 \rangle}_{\subset G_0} \coprod \underbrace{T_n \left\langle \alpha_0^2 \right\rangle}_{\subset G_0} \coprod \underbrace{T_n \alpha_0 \left\langle \alpha_0^2 \right\rangle}_{\subset G_2}.$$

The inclusions follow from the fact that the elements of $T_n \langle \alpha_0^2 \rangle$ (resp. $T_n\alpha_0 \langle \alpha_0^2 \rangle$) are conjugate (in Dic_{4n}), $T_n \in G_0$, and

$$\begin{aligned}\pi(T_n\alpha_0) &= (1, n)(2, n-1) \cdots \left(\frac{n}{2}, \frac{n}{2}+1\right)(1, n, \ldots, 2)\\ &= (n)\left(\frac{n}{2}\right)(1, n-1)(2, n-2)(3, n-3) \cdots \left(\frac{n}{2}-1, \frac{n}{2}+1\right),\end{aligned}$$

where $\pi\colon B_n(\mathbb{S}^2) \longrightarrow S_n$ denotes the homomorphism defined on the generators by $\pi(\sigma_i) = (i, i+1)$. Thus $T_n\alpha_0 \in G_2$.

If $k \nmid \frac{n}{2}$ then by Proposition 5, there is just one conjugacy class of Dic_{4k} of the form $\left\langle \alpha_0^{n/k}, T_n \right\rangle$, and since n/k is odd, we have

$$\mathrm{Dic}_{4k} = \underbrace{\left\langle \alpha_0^{n/k} \right\rangle}_{\subset G_0} \coprod \underbrace{T_n \left\langle \alpha_0^{n/k} \right\rangle}_{\subset G_2}.$$

In particular, all of the elements of Dic_{4k} of order 4 belong to G_2. Thus $\mathrm{Dic}_{4k} \cap (G_0 \backslash G_2) \neq \varnothing$ and $\mathrm{Dic}_{4k} \cap (G_2 \backslash G_0) \neq \varnothing$.

If $k \mid \frac{n}{2}$ then by Proposition 5, there are two non-conjugate copies of Dic_{4k} given by

$$\left\langle \alpha_0^{n/k}, T_n \right\rangle = \underbrace{\left\langle \alpha_0^{n/k} \right\rangle}_{\subset G_0} \coprod \underbrace{T_n \left\langle \alpha_0^{n/k} \right\rangle}_{\subset G_0}, \text{ and}$$

and

$$\left\langle \alpha_0^{n/k}, T_n\alpha_0 \right\rangle = \underbrace{\left\langle \alpha_0^{n/k} \right\rangle}_{\subset G_0} \coprod \underbrace{T_n\alpha_0 \left\langle \alpha_0^{n/k} \right\rangle}_{\subset G_2}.$$

The first copy lies entirely within G_0, while the second lies in $G_0 \cup G_2$ and meets both $G_0 \backslash G_2$ and $G_2 \backslash G_0$.

A similar result holds for $\mathrm{Dic}_{4(n-2)}$: its subgroups are either subgroups of $\mathbb{Z}_{2(n-2)}$, so lie in G_2, or else are dicyclic, of the form Dic_{4k}, where $k \mid n-2$. If $k = 1$ then the subgroup in question is $\langle T_n \rangle$ which lies in G_0. If $k > 1$ then as above, we distinguish two cases. If $k \nmid \frac{n-2}{2}$ then there is just one copy of Dic_{4k} which lies in $G_0 \cup G_2$ and meets both $G_0 \backslash G_2$ and $G_2 \backslash G_0$. If $k \mid \frac{n-2}{2}$, then setting $\alpha_2' = \alpha_0 \alpha_2 \alpha_0^{-1}$, there are two copies of Dic_{4k}, $\left\langle \alpha_2'^{n/k}, T_n \right\rangle$, which lies in $G_0 \cup G_2$ and meets both $G_0 \backslash G_2$ and $G_2 \backslash G_0$, and $\left\langle \alpha_2'^{n/k}, \alpha_2' T_n \right\rangle$, which is contained in G_2.
(b) Suppose that H is a subgroup of a copy of T_1 when T_1 is maximal, so $n \equiv 4 \bmod 6$. Assume first that $H \cong T_1$. Since $H \cong \mathcal{Q}_8 \rtimes \mathbb{Z}_3$, all of its order 4 elements are conjugate, and so all elements of $\mathcal{Q}_8$ must lie in the same G_i. Now $\mathcal{Q}_8 = \mathrm{Dic}_8$, so from above, we must be in one of the cases $2 \mid \frac{n}{2}$ or $2 \mid \frac{n-2}{2}$. Indeed if $n \equiv 4 \bmod 12$ then $n = 4 + 12l = 4(1 + 3l)$, $l \in \mathbb{N}$, and so $\mathcal{Q}_8$ is contained in G_0, while if $n \equiv 10 \bmod 12$ then $n = 10 + 12l = 2(5 + 6l)$, $l \in \mathbb{N}$, and so $\mathcal{Q}_8$ is contained in G_2. The remaining elements of H are of order 3 or 6, and since $n \equiv 4 \bmod 6$, lie in G_1. So if $n \equiv 4 \bmod 12$ (resp. $n \equiv 10 \bmod 12$) then H lies in $G_0 \cup G_1$ (resp. $G_2 \cup G_1$) and meets both G_0 (resp. G_2) and G_1.

From this, we deduce immediately the following: if H is isomorphic to $\mathbb{Z}_3$ or $\mathbb{Z}_6$ then it is contained in G_1, and if it is isomorphic to $\mathbb{Z}_4$ or $\mathcal{Q}_8$ then it is contained in G_0 if $n \equiv 4 \bmod 12$, and in G_2 if $n \equiv 10 \bmod 12$.
(c) Suppose that H is a subgroup of a copy of I when I is maximal, so $n \equiv 0, 2, 12, 20 \bmod 30$. Assume first that $H \cong I$. So I has a subgroup isomorphic to T_1, whose copy of $\mathcal{Q}_8$ lies entirely in G_0 or G_2. The subgroups of order 8 of H are its Sylow 2-subgroups, so are conjugate, and thus all lie either in G_0 or in G_2. Hence from the analysis of the dicyclic case, 2 divides $\frac{n}{2}$ or $\frac{n-2}{2}$. Further, all elements of H of order 4 are contained in one of its subgroups isomorphic to $\mathcal{Q}_8$ (because the order 2 elements of A_5 are the product of two transpositions, and are contained in a subgroup isomorphic to $\mathbb{Z}_2 \oplus \mathbb{Z}_2$, which lifts to $\mathcal{Q}_8$ in I). Hence all order 4 elements of H lie either in G_0 if $4 \mid n$, or in G_2 if $4 \mid n-2$. The remaining elements of H are of order $3, 6, 5$ and 10, and lie in either G_0 or G_2 depending on the value of n modulo the order. Thus H lies entirely in G_0 (resp. G_2) if $n \equiv 0 \bmod 60$ (resp. $n \equiv 2 \bmod 60$), and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 12, 20, 30, 32, 42, 50 \bmod 60$.

We now consider the other possibilities for subgroups of I: if H is isomorphic to either $\mathbb{Z}_3$ or $\mathbb{Z}_6$, it is contained in G_0 if $n \equiv 0, 12 \bmod 30$, and in G_2 if $n \equiv 2, 20 \bmod 30$; if H is isomorphic to either $\mathbb{Z}_5$ or $\mathbb{Z}_{10}$, it is contained in G_0 if $n \equiv 0, 20 \bmod 30$, and in G_2 if $n \equiv 2, 12 \bmod 30$; and if H is isomorphic to either $\mathbb{Z}_4$ or $\mathcal{Q}_8$, it is contained in G_0 if $n \equiv 0, 12, 20, 32 \bmod 60$, and in G_2 if $n \equiv 2, 30, 42, 50 \bmod 60$. Next, if H is isomorphic to T_1, it consists of a copy of $\mathcal{Q}_8$ and elements of order 3 and 6, so lies in G_0 if $n \equiv 0, 12 \bmod 60$, in G_2 if $n \equiv 2, 50 \bmod 60$, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 20, 30, 32, 42 \bmod 60$. Now suppose that H is isomorphic to $\mathrm{Dic}_{12} \cong \mathbb{Z}_3 \rtimes \mathbb{Z}_4 = \mathbb{Z}_6 \coprod T_n \mathbb{Z}_6$. Since the elements of $T_n \mathbb{Z}_6$ are of order 4, it follows from the analysis of the cyclic subgroups that H satisfies the same conditions as in the case of T_1. Finally, if H is isomorphic to $\mathrm{Dic}_{20} \cong \mathbb{Z}_5 \rtimes \mathbb{Z}_4 = \mathbb{Z}_{10} \coprod T_n \mathbb{Z}_{10}$, since the elements of $T_n \mathbb{Z}_{10}$ are of order 4, it follows from the analysis of the cyclic subgroups that H lies in G_0 if $n \equiv 0, 20 \bmod 60$, in G_2 if $n \equiv 2, 42 \bmod 60$, and lies in $G_0 \cup G_2$ and meets both

G_0 and G_2 if $n \equiv 12, 30, 32, 50 \bmod 60$.

(d) Suppose that H is a subgroup of a copy of O_1 when O_1 is maximal, so $n \equiv 0, 2 \bmod 6$. Assume first that $H \cong O_1$. Then it has a subgroup isomorphic to T_1 (which is unique since S_4 has a unique subgroup abstractly isomorphic to A_4), and the copy of $\mathcal{Q}_8$ lying in T_1 lies entirely in G_0 if $n \equiv 0, 8 \bmod 12$, and in G_2 if $n \equiv 2, 6 \bmod 12$. The complement of this copy of $\mathcal{Q}_8$ in T_1 consists of elements of order 3 and 6, and so lie in G_0 if $n \equiv 0 \bmod 6$ and in G_2 if $n \equiv 2 \bmod 6$ (thus the subgroups of O_1 isomorphic to $\mathbb{Z}_3$ and $\mathbb{Z}_6$ lie in G_0 if $n \equiv 0 \bmod 6$ and in G_2 if $n \equiv 2 \bmod 6$). Thus T_1 lies in G_0 if $n \equiv 0 \bmod 12$, in G_2 if $n \equiv 2 \bmod 12$, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 6, 8 \bmod 12$.

In order to analyse the remaining possible subgroups $\mathcal{Q}_{16}$, Dic_{12}, Dic_{20} of O_1, as well as the other copy of $\mathcal{Q}_8$ lying in $\mathcal{Q}_{16}$, we must study the elements of $H \backslash T_1$. They project to elements of $S_4 \backslash A_4$, which are either 4-cycles, or transpositions. We analyse the geometric formulation of O_1 described in Section 3 as being obtained from the action of S_4 on a cube, with the n strings attached appropriately. The 4-cycles are realised by rotations by $\pi/2$ about an axis which passes through the centres of two opposite faces. This gives rise to an element of G_0 if the n marked points are not these central points (i.e. if $n \equiv 0, 8 \bmod 12$), and to elements of G_2 if some of the n marked points are central points of the faces (i.e. if $n \equiv 2, 6 \bmod 12$). The transpositions are realised by rotations by π about an axis which passes through the centres of two diagonally-opposite edges. This gives rise to an element of G_0 if there are an even number of marked points on each edge (i.e. if $n \equiv 0, 6, 8, 14 \bmod 24$), and to elements of G_2 if there are an odd number of marked points on each edge (i.e. if $n \equiv 2, 12, 18, 20 \bmod 24$). Putting together these results with those for T_1, if $H \cong O_1$, we conclude that it lies in G_0 if $n \equiv 0 \bmod 24$, in G_2 if $n \equiv 2 \bmod 24$, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 6, 8, 12, 14, 18, 20 \bmod 24$.

Now suppose that H is a subgroup of a copy of O_1 isomorphic to $\mathcal{Q}_{16}$. Such subgroups are the Sylow 2-subgroups of O_1, so are conjugate. If $n \equiv 0 \bmod 24$ (resp. $n \equiv 2 \bmod 24$) then O_1 lies in G_0 (resp. G_2), and hence so does $\mathcal{Q}_{16}$. So suppose that $n \not\equiv 0, 2 \bmod 24$. Any subgroup of O_1 isomorphic to $\mathcal{Q}_{16}$ contains elements of order 8 which lie in $O_1 \backslash T_1$, and so are associated with the above 4-cycles. Further, H projects to a subgroup of S_4 isomorphic to D_8 which is generated by a 4-cycle and a transposition. Studying the associated rotations as above, if one has fixed points and the other not then automatically H lies in $G_0 \cup G_2$ and meets both G_0 and G_2. This occurs when $n \equiv 6, 12, 14, 20 \bmod 24$. So suppose that $n \equiv 8, 18 \bmod 24$.

If $n \equiv 8 \bmod 24$ (resp. $n \equiv 18 \bmod 24$) then the elements of H corresponding to the 4-cycles and the transpositions of D_8 belong to G_0 (resp. G_2). Further, the remaining elements of D_8 are products of such elements, and so the corresponding elements in H are also elements of $T_1 \cong \mathcal{Q}_8 \rtimes \mathbb{Z}_3$ of order 4. But such elements lie in the $\mathcal{Q}_8$-factor. Since $n \equiv 8 \bmod 12$ (resp. $n \equiv 6 \bmod 12$), this copy of $\mathcal{Q}_8$ lies in G_0 (resp. G_2), and hence so does the given subgroup $\mathcal{Q}_{16}$. Summing up, H lies in G_0 if $n \equiv 0, 8 \bmod 24$, in G_2 if $n \equiv 2, 18 \bmod 24$, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 6, 12, 14, 20 \bmod 24$.

Now suppose that H is a subgroup of a copy of O_1 isomorphic to Dic_{12}. If $n \equiv 0 \bmod 24$ (resp. $n \equiv 2 \bmod 24$) then O_1 lies in G_0 (resp. G_2), and hence so does H. So suppose that $n \not\equiv 0, 2 \bmod 24$. Any subgroup of O_1 isomorphic to H projects onto a subgroup of S_4 isomorphic to S_3 which consists of 3-cycles and transpositions. Hence H is generated by an element of order 4 lying in $O_1 \backslash T_1$, and an element of order 6, which lies in T_1. The first element belongs to G_0 if $n \equiv 6, 8, 14 \bmod 24$ and to G_2 if $n \equiv 12, 18, 20 \bmod 24$, while the second element belongs to G_0 if $n \equiv 6, 12, 18 \bmod 24$ and to G_2 if $n \equiv 8, 14, 20 \bmod 24$. Hence if $n \equiv 8, 12, 14, 18 \equiv 24$ then H lies in $G_0 \cup G_2$ and meets both G_0 and G_2. The product of the two given generators is also of order 4 and so lies in G_0 if $n \equiv 6 \bmod 24$, and in G_2 if $n \equiv 20 \bmod 24$. Thus H lies in G_0 if $n \equiv 0, 6 \bmod 24$, in G_2 if $n \equiv 2, 20 \bmod 24$, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 8, 12, 14, 18 \bmod 24$.

Now suppose that H is a subgroup of a copy of O_1 isomorphic to $\mathbb{Z}_4$. There are two possibilities. If it is contained in the copy of $\mathcal{Q}_8$ lying in the subgroup T_1, from the results for $\mathcal{Q}_8$, we see that H lies in G_0 if $n \equiv 0, 8 \bmod 12$, and in G_2 if $n \equiv 2, 6 \bmod 12$. The second possibility is that H possesses elements in $O_1 \backslash T_1$, and emanates from the rotation of order 2 whose permutation is a transposition. Thus it is contained in G_0 if $n \equiv 0, 6, 8, 14 \bmod 24$, and to G_2 if $n \equiv 2, 12, 18, 20 \bmod 24$.

Finally, suppose that H is a subgroup of a copy of O_1 isomorphic to $\mathcal{Q}_8$. Again there are two possibilities. If H lies in the subgroup T_1, it is contained in G_0 if $n \equiv 0, 8 \bmod 12$, and to G_2 if $n \equiv 2, 6 \bmod 12$. The second possibility is that it projects to a subgroup of S_4 generated by two transpositions having disjoint support. Such a subgroup thus has four elements of order 4 in $O_1 \backslash T_1$ and two in T_1. From the results obtained in the case of $\mathbb{Z}_4$, we see that H lies in G_0 if $n \equiv 0, 8 \bmod 24$, in G_2 if $n \equiv 2, 18 \bmod 24$, and lies in $G_0 \cup G_2$ and meets both G_0 and G_2 if $n \equiv 6, 12, 14, 20 \bmod 24$.

$\square$

5 Realisation of finite groups as subgroups of the lower central and derived series of $B_n(\mathbb{S}^2)$

In this section, we consider the realisation of the finite subgroups of Theorem 3 as subgroups of the lower central $\Gamma_i(B_n(\mathbb{S}^2))$ and derived series $(B_n(\mathbb{S}^2)^{(i)})$ of $B_n(\mathbb{S}^2)$. By [GG4], we already know that if $4 \mid n$ then $\Gamma_2(B_n(\mathbb{S}^2))$ has a subgroup isomorphic to $\mathcal{Q}_8$. If $n \geqslant 4$ is even but not divisible by 4, we may ask if the same result is true if $4 \nmid n$. We start by proving Theorem 6, which is the case of the dicyclic groups. We then then complete the analysis of the other finite subgroups in Proposition 16.

Proof of Theorem 6. Suppose that n is even. Let $N \in \{n-2, n\}$, set $N = 2^l k$ where $l \in \mathbb{N}$ and k is odd, and let $x = \alpha_0$ (resp. $x = \alpha_0\alpha_2\alpha_0^{-1}$) if $N = n$ (resp. $N = n-2$).

(a) Since $B_n(\mathbb{S}^2)$ has a subgroup $\langle x, T_n \rangle$ isomorphic to $\mathrm{Dic}_{4N} = \mathrm{Dic}_{2^{l+2}k}$, the statement is true for $j = 0$. So suppose the result holds for some $j \in \{0, 1, \ldots, l-1\}$. Then $B_n(\mathbb{S}^2)$ contains 2^j copies of $\mathrm{Dic}_{2^{l+2-j}k}$ of the form $\left\langle x^{2^j}, x^i T_n \right\rangle$, for $i = 0, 1, \ldots, 2^j - 1$. Hence $\left\langle x^{2^{j+1}}, x^i T_n \right\rangle$ is a subgroup of $\left\langle x^{2^j}, x^i T_n \right\rangle$ isomorphic to $\mathrm{Dic}_{2^{l+1-j}k}$. But since

$$\left(x^{(2^j+i)}T_n\right)^2 = x^{(2^j+i)}T_n x^{(2^j+i)}T_n^{-1}T_n^2 = \Delta_n, \quad \text{and}$$
$$x^{(2^j+i)}T_n \cdot x^{2^{j+1}} \left(x^{(2^j+i)}T_n\right)^{-1} = x^{-2^{j+1}},$$

it follows that $\left\langle x^{2^{j+1}}, x^{(2^j+i)}T_n \right\rangle$ is also a subgroup of $\left\langle x^{2^j}, x^i T_n \right\rangle$ isomorphic to $\mathrm{Dic}_{2^{l+1-j}k}$.

If q is any divisor of k, then replacing x by x^q yields also 2^j copies $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle$, $i = 0, 1, \ldots, 2^j - 1$, of $\mathrm{Dic}_{2^{l+2-j}k/q}$ for $j \in \{0, 1, \ldots, l\}$.

(b) If $j = 0$, the statement holds trivially. So suppose that $j \geqslant 1$. From part (a), $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle$ and $\left\langle x^{2^j q}, x^{i'q} T_n \right\rangle$ are subgroups of $B_n(\mathbb{S}^2)$ isomorphic to $\mathrm{Dic}_{2^{l+2-j}k/q}$. Under the Abelianisation homomorphism $\xi\colon B_n(\mathbb{S}^2) \longrightarrow \mathbb{Z}_{2(n-1)}$, $\xi(x) = \overline{n-1}$, and

$$\xi(T_n) = \xi\big((\sigma_1 \cdots \sigma_{n-1}) \cdots (\sigma_1\sigma_2)\sigma_1\big) = \overline{\frac{1}{2}n(n-1)} = \begin{cases} \overline{0} & \text{if } \frac{n}{2} \text{ is even} \\ \overline{n-1} & \text{if } \frac{n}{2} \text{ is odd.} \end{cases}$$

Since $j \geqslant 1$, $\xi(x^{2^j q}) = \overline{0}$. Furthermore,

$$\xi\left(x^{iq} T_n\right) = \begin{cases} \overline{0} & \text{if } \frac{n}{2} + i \text{ is even} \\ \overline{n-1} & \text{if } \frac{n}{2} + i \text{ is odd.} \end{cases}$$

So $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle \subset \Gamma_2(B_n(\mathbb{S}^2))$ if and only if $\frac{n}{2} + i$ is even. Thus if $i - i'$ is odd, the subgroups $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle$ and $\left\langle x^{2^j q}, x^{i'q} T_n \right\rangle$ cannot be conjugate. But by Proposition 5(b), these are precisely the conjugacy classes of subgroups isomorphic to $\mathrm{Dic}_{2^{l+2-j}k/q}$. The result follows.

□

From this, we may deduce Proposition 7.

Proof of Proposition 7. We use the notation of the proof of Theorem 6. If $j = 0$ and q is an odd divisor of n then there is just one conjugacy class of the abstract group $\mathrm{Dic}_{4n/q}$, which is realised as $\langle x^q, T_n \rangle$. Now $x^q \notin \Gamma_2(B_n(\mathbb{S}^2))$, so $\mathrm{Dic}_{4n/q} \not\subset \Gamma_2(B_n(\mathbb{S}^2))$.

If $j \geqslant 1$ then as we saw in the proof of Theorem 6, $\left\langle x^{2^j q}, x^{iq} T_n \right\rangle \subset \Gamma_2(B_n(\mathbb{S}^2))$ if and only if $\frac{n}{2} + i$ is even. So with $i = 0, 1$, one of $\left\langle x^{2^j q}, T_n \right\rangle$ and $\left\langle x^{2^j q}, x^q T_n \right\rangle$ is contained in $\Gamma_2(B_n(\mathbb{S}^2))$, while the other is not.

Finally, let N be the element of $\{n, n-2\}$ divisible by 4. Then $l \geqslant 2$, and taking $q = k$ and $j = l - 1$, from the previous paragraph, one of $\left\langle x^{N/2}, T_n \right\rangle$ and $\left\langle x^{N/2}, x^k T_n \right\rangle$ (the two non-conjugate copies of Q_8) belongs to $\Gamma_2(B_n(\mathbb{S}^2))$, the other not. □

We now give the analogous result for the cyclic and binary polyhedral subgroups of $B_n(\mathbb{S}^2)$.

PROPOSITION 16. *Let G be a finite subgroup of $B_n(\mathbb{S}^2)$.*

(a) Suppose that G is cyclic.

(i) If G is of order 2, then $G \subset \Gamma_2(B_n(\mathbb{S}^2))$ if and only if n is even.

(ii) Suppose that G is of order greater than or equal to 3. Then either:

– *$|G|$ divides $2(n-1)$ in which case $G \not\subset \Gamma_2(B_n(\mathbb{S}^2))$, or*

– *$|G|$ divides $2(n-i)$, where $i \in \{0, 2\}$. In this case, $G \subset \Gamma_2(B_n(\mathbb{S}^2))$ if and only if $|G|$ divides $n - i$.*

(b) Suppose that G is a subgroup of order at least 3 of some binary polyhedral subgroup H of $B_n(\mathbb{S}^2)$.

(i) Suppose that $H \cong T_1$ in the case that T_1 is maximal. Then $G \subset \Gamma_2(B_n(\mathbb{S}^2))$ if $G \cong \mathbb{Z}_4, \mathcal{Q}_8$, and $G \not\subset \Gamma_2(B_n(\mathbb{S}^2))$ if $G \cong \mathbb{Z}_3, \mathbb{Z}_6, T_1$.

(ii) Suppose that $H \cong I$ in the case that I is maximal. Then $G \subset \Gamma_2(B_n(\mathbb{S}^2))$.

(iii) Suppose that $H \cong O_1$ in the case that O_1 is maximal. If G is contained in the subgroup K of H isomorphic to T_1 then $G \subset \Gamma_2(B_n(\mathbb{S}^2))$. If $G \not\subset K$ then $G \subset \Gamma_2(B_n(\mathbb{S}^2))$ if $n \equiv 0, 2, 8, 18 \bmod 24$, and $G \not\subset \Gamma_2(B_n(\mathbb{S}^2))$ if $n \equiv 6, 12, 14, 20 \bmod 24$.

Proof. We set $\Gamma_2 = \Gamma_2(B_n(\mathbb{S}^2))$. If G is of order 2, then $G = \langle \Delta_n \rangle$ and as $\xi(\Delta_n) = \overline{n(n-1)}$, it follows easily that $G \subset \Gamma_2$ if and only if n is even. We assume from now on that $|G| \geqslant 3$. Since Γ_2 is normal in $B_n(\mathbb{S}^2)$, we may work up to conjugation.

First suppose that G is cyclic. Then by Theorem 1, it is conjugate to a subgroup of $\langle \alpha_i \rangle$ for some $i \in \{0, 1, 2\}$. If $i = 1$ then $\xi(\alpha_1^j) = \overline{jn}$ for all $j \in \mathbb{Z}$. If $\alpha_1^j \in \Gamma_2$ then there exists $k \in \mathbb{Z}$ such that $jn = 2k(n-1)$, thus $n - 1 \mid j$, and so $j = l(n-1)$ for some $l \in \mathbb{Z}$. But then $\alpha_1^j = \alpha_1^{l(n-1)} \in \langle \Delta_n \rangle$. We conclude that $\langle \alpha_1 \rangle \cap \Gamma_2 \subset \langle \Delta_n \rangle$. Hence $G \not\subset \Gamma_2$.

Suppose then that G is conjugate to a subgroup of $\langle \alpha_i \rangle$, where $i = 0, 2$. Set $k = |G|$. Then $\xi(\alpha_i) = \overline{n-1}$, $k \mid 2(n-i)$, and up to conjugacy, $G = \left\langle \alpha_i^{2(n-i)/k} \right\rangle$. So $G \subset \Gamma_2$ if and only if $2(n-i)/k$ is even, which is equivalent to $k \mid n - i$. Thus if G is conjugate to a subgroup of $\langle \alpha_i \rangle$, where $i = 0, 2$, we have:

$$G \subset \Gamma_2 \Longleftrightarrow |G| \mid n - i. \tag{6}$$

Now suppose that H is isomorphic to T_1 in the case that T_1 is maximal, so that $n \equiv 4 \bmod 6$. If G is isomorphic to $T_1, \mathbb{Z}_6$ or $\mathbb{Z}_3$ then the order 3 elements lie in $G_1 \backslash \langle \Delta_n \rangle$, and from the cyclic case, it follows that $G \not\subset \Gamma_2$. So assume that G is isomorphic to either $\mathbb{Z}_4$ or $\mathcal{Q}_8$. Since $\mathcal{Q}_8$ is generated by elements of order 4, it suffices to analyse the case $\mathbb{Z}_4$. By Proposition 15, G lies in G_0 if $n \equiv 4 \bmod 12$, and in G_2 if $n \equiv 10 \bmod 12$. In both cases, $G \subset \Gamma_2$ by equation (6).

Now suppose that H is isomorphic to I in the case that I is maximal, so that $n \equiv 0, 2, 12, 20 \bmod 30$. We claim that $G \subset \Gamma_2$ whatever the value of n. To see this, it suffices to check that all of the maximal cyclic subgroups $\mathbb{Z}_4$, $\mathbb{Z}_6$, $\mathbb{Z}_{10}$ of I are contained in Γ_2. This follows easily from Proposition 15 and equation (6).

Now suppose that H is isomorphic to O_1 in the case that O_1 is maximal, so that $n \equiv 0, 2 \bmod 6$. Again it suffices to consider the maximal cyclic subgroups $\mathbb{Z}_4$, $\mathbb{Z}_6$ and $\mathbb{Z}_8$ of O_1. Applying Proposition 15 and equation (6), we obtain the following results:

- if G is isomorphic to $\mathbb{Z}_8$, it projects to a subgroup of S_4 generated by a 4-cycle. Then $G \subset G_0$ if $n \equiv 0, 8 \bmod 12$, and $G \subset G_2$ if $n \equiv 2, 6 \bmod 12$, and so $G \subset \Gamma_2$ if $n \equiv 0, 2, 8, 18 \bmod 24$, and $G \not\subset \Gamma_2$ if $n \equiv 6, 12, 14, 20 \bmod 24$.
- if G is isomorphic to $\mathbb{Z}_6$ then $G \subset \Gamma_2$.
- if G is isomorphic to $\mathbb{Z}_4$, there are two possibilities. If G lies in the subgroup K of O_1 isomorphic to T_1 then $G \subset \Gamma_2$. Otherwise G is generated by an element of order 4 not belonging to K, in which case we obtain the same answer as for $\mathbb{Z}_8$.

Since every cyclic subgroup of order 3 of O_1 is contained in one of order 6, this gives the results if G is cyclic. Suppose now that $G = K$. Then G is generated by the elements of order 6 and the elements of order 4 belonging to K, so $G \subset \Gamma_2$.

If G is abstractly isomorphic to $\mathcal{Q}_{16}$ then it is generated by elements of order 8, elements of order 4 lying in K, and elements of order 4 not lying in K. From above, we have that $G \subset \Gamma_2$ if $n \equiv 0, 2, 8, 18 \bmod 24$, and $G \not\subset \Gamma_2$ if $n \equiv 6, 12, 14, 20 \bmod 24$.

If G is abstractly isomorphic to $\mathcal{Q}_8$ then there are two possibilities: either G lies in K, so is contained in Γ_2, or else it is generated by elements of order 4 not belonging to K. In this case, from above, $G \subset \Gamma_2$ if $n \equiv 0, 2, 8, 18 \bmod 24$, and $G \not\subset \Gamma_2$ if $n \equiv 6, 12, 14, 20 \bmod 24$.

Finally, suppose that G is abstractly isomorphic to Dic_{12}. Then it projects to a copy of S_3 in S_4. From above, it follows that $G \subset \Gamma_2$ if $n \equiv 0, 2, 8, 18 \bmod 24$, and $G \not\subset \Gamma_2$ if $n \equiv 6, 12, 14, 20 \bmod 24$. ☐

REMARK 17. Having dealt with the behaviour of the finite subgroups relative to the commutator subgroup of $B_n(\mathbb{S}^2)$, one might ask what happens for the higher elements of the lower central series $\left\{\Gamma_i(B_n(\mathbb{S}^2))\right\}_{n\in\mathbb{N}}$ and of the derived series $\left\{\left(B_n(\mathbb{S}^2)\right)^{(i)}\right\}_{i\geqslant 0}$ of $B_n(\mathbb{S}^2)$. But if $n \neq 2$ (resp. $n \geqslant 5$), the lower central series (resp. derived series) of $B_n(\mathbb{S}^2)$ is stationary from the commutator subgroup onwards [GG3]. It just remains to look at the derived series of $B_4(\mathbb{S}^2)$. Recall from that paper that $(B_4(\mathbb{S}^2))^{(1)}$ is a semi-direct product of $\mathcal{Q}_8$ by a free group $\mathbb{F}_2$ of rank two, that $(B_4(\mathbb{S}^2))^{(2)}$ is a semi-direct product of $\mathcal{Q}_8$ by the derived subgroup $(\mathbb{F}_2)^{(1)}$ of $\mathbb{F}_2$, that $(B_4(\mathbb{S}^2))^{(3)}$ is the direct product of $\langle \Delta_4 \rangle$ by $(\mathbb{F}_2)^{(2)}$, and that $(B_4(\mathbb{S}^2))^{(i+1)} \cong (\mathbb{F}_2)^{(i)}$ for all $i \geqslant 3$. Thus there is a copy of $\mathcal{Q}_8$ which lies in $(B_4(\mathbb{S}^2))^{(2)}$ but not in $(B_4(\mathbb{S}^2))^{(3)}$. The full twist remains until $(B_4(\mathbb{S}^2))^{(3)}$, and then $(B_4(\mathbb{S}^2))^{(4)}$ is torsion free.

6 A proof of Murasugi's theorem

Let H_1, H_2 be isomorphic finite cyclic subgroups of $\mathcal{M}_{0,n}$. From Theorem 9, if n is odd, or if n is even and $|H_1| = |H_2| \neq 2$ then H_1 and H_2 are conjugate. If n is even, there are exactly two conjugacy classes of subgroups of $\mathcal{M}_{0,n}$ of order 2, and thus there are exactly two conjugacy classes of subgroups of $B_n(\mathbb{S}^2)$ of order 4.

It follows from Section 2 that:

PROPOSITION 18. *Let G_1, G_2 be isomorphic finite cyclic subgroups of order m of $B_n(\mathbb{S}^2)$. If n is odd, or if n is even and $m \neq 4$ then G_1 and G_2 are conjugate. If n is even, there are exactly two conjugacy classes of subgroups of $B_n(\mathbb{S}^2)$ of order* 4. ☐

If n is even then $\alpha_0^{n/2}$ and $\alpha_2^{(n-2)/2}$ are of order 4, and they generate non-conjugate subgroups since their images in S_n are not conjugate, which yields the two conjugacy classes of $\mathbb{Z}_4$ of Proposition 18. From this, we may deduce Theorem 1.

Proof of Theorem 1. Let $x \in B_n(\mathbb{S}^2)$ be a torsion element. Then $\langle x \rangle$ is contained in a maximal cyclic subgroup C of one of the maximal finite subgroups G of $B_n(\mathbb{S}^2)$ given by Theorem 3.

First suppose that n is odd. Then G is one of $\mathbb{Z}_{2(n-1)}$, Dic_{4n} and $\mathrm{Dic}_{4(n-2)}$, and so C must be one of $\mathbb{Z}_{2(n-1)}$, $\mathbb{Z}_{2n}$, $\mathbb{Z}_{2(n-2)}$ and $\mathbb{Z}_4$. Hence C is isomorphic to $\langle \alpha_1 \rangle$, $\langle \alpha_0 \rangle$, $\langle \alpha_2 \rangle$ and $\left\langle \alpha_1^{(n-1)/2} \right\rangle$ respectively. So by Proposition 18, x is conjugate to a power of one of α_0, α_1 and α_2 which proves the theorem in this case.

Now suppose that n is even. If $C \cong \mathbb{Z}_4$ then C is conjugate to one of $\left\langle \alpha_0^{n/2} \right\rangle$ or $\left\langle \alpha_2^{(n-2)/2} \right\rangle$, and the result holds. So suppose that $C \ncong \mathbb{Z}_4$. If G is one of $\mathbb{Z}_{2(n-1)}$, Dic_{4n} and $\mathrm{Dic}_{4(n-2)}$, then C is one of $\mathbb{Z}_{2(n-1)}$, $\mathbb{Z}_{2n}$, $\mathbb{Z}_{2(n-2)}$, and so is isomorphic to $\langle \alpha_1 \rangle$, $\langle \alpha_0 \rangle$, and $\langle \alpha_2 \rangle$ respectively. If $G = T_1$ (so $n \equiv 4 \bmod 6$) then $C \cong \mathbb{Z}_6$, and so is conjugate to $\left\langle \alpha_1^{(n-1)/3} \right\rangle$. If $G = O_1$ (so $n \equiv 0, 2 \bmod 6$) then $C \cong \mathbb{Z}_6$ or $C \cong \mathbb{Z}_8$, and so is conjugate to $\left\langle \alpha_0^{n/3} \right\rangle$ or $\left\langle \alpha_2^{(n-2)/3} \right\rangle$. Finally, if $G = I$ (so $n \equiv 0, 2, 12, 20 \bmod 30$) then C is isomorphic to one of $\mathbb{Z}_6$ or $\mathbb{Z}_{10}$. If $C \cong \mathbb{Z}_6$ then C is conjugate to $\left\langle \alpha_0^{n/3} \right\rangle$ if $n \equiv 0, 12 \bmod 30$ or to $\left\langle \alpha_2^{(n-2)/3} \right\rangle$ if $n \equiv 2, 20 \bmod 30$. If $C \cong \mathbb{Z}_{10}$ then C is conjugate to $\left\langle \alpha_0^{n/5} \right\rangle$ if $n \equiv 0, 20 \bmod 30$ or to $\left\langle \alpha_2^{(n-2)/5} \right\rangle$ if $n \equiv 2, 12 \bmod 30$. In all cases, x is conjugate to a power of one of α_0, α_1 and α_2, which completes the proof of the theorem. $\square$

References

[AM] A. Adem and R. J. Milgram, Cohomology of finite groups, Springer-Verlag, New York-Heidelberg-Berlin (1994).

[ATD] Algebraic topology discussion list, January 2004, `http://www.lehigh.edu/~dmd1/pz119.txt`.

[A1] E. Artin, Theorie der Zöpfe, *Abh. Math. Sem. Univ. Hamburg* **4** (1925), 47–72.

[A2] E. Artin, Theory of braids, *Ann. Math.* **48** (1947), 101–126.

[Bi] J. S. Birman, Braids, links and mapping class groups, *Ann. Math. Stud.* **82**, Princeton University Press, 1974.

[Br] K. S. Brown, Cohomology of groups, Graduate Texts in Mathematics **87**, Springer-Verlag, New York-Berlin (1982).

[CG] F. R. Cohen and S. Gitler, On loop spaces of configuration spaces, *Trans. Amer. Math. Soc.* **354** (2002), 1705–1748.

[Co] H. S. M. Coxeter, Regular complex polytopes, Second edition, Cambridge University Press, Cambridge, 1991.

[CM] H. S. M. Coxeter and W. O. J. Moser, Generators and relations for discrete groups, Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 14, Fourth edition, Springer-Verlag, Berlin, 1980.

[EE] C. J. Earle and J. Eells, The diffeomorphism group of a compact Riemann surface, *Bull. Amer. Math. Soc.* **73** (1967), 557–559.

[F] E. Fadell, Homotopy groups of configuration spaces and the string problem of Dirac, *Duke Math. Journal* **29** (1962), 231–242.

[FH] E. Fadell and S. Y. Husseini, Geometry and topology of configuration spaces, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2001.

[FVB] E. Fadell and J. Van Buskirk, The braid groups of $\mathbb{E}^2$ and $\mathbb{S}^2$, *Duke Math. Journal* **29** (1962), 243–257.

[FoN] R. H. Fox and L. Neuwirth, The braid groups, *Math. Scandinavica* **10** (1962), 119–126.

[GVB] R. Gillette and J. Van Buskirk, The word problem and consequences for the braid groups and mapping class groups of the 2-sphere, *Trans. Amer. Math. Soc.* **131** (1968), 277–296.

[GoG1] M. Golasiński and D. L. Gonçalves, Spherical space forms – homotopy types and self-equivalences, The Skye Conference Proceedings, *Progr. Math.* **215** Birkhäuser, Basel (2004), 153–165.

[GoG2] M. Golasiński and D. L. Gonçalves, Spherical space forms – homotopy types and self-equivalences for the groups $\mathbb{Z}/a \rtimes \mathbb{Z}/b$ and $\mathbb{Z}/a \rtimes (\mathbb{Z}/b \times \mathcal{Q}_{2i})$, *Topology Appl.* **146–147** (2005), 451–470.

[GoG3] M. Golasiński and D. L. Gonçalves, Spherical space forms – homotopy types and self-equivalences for the groups $\mathbb{Z}/a \rtimes (\mathbb{Z}/b \times T_i^\star)$ and $\mathbb{Z}/a \rtimes (\mathbb{Z}/b \times O_n^\star)$, *J. Homotopy and Related Structures* **1** (2006), 29–45.

[GoG4] M. Golasiński and D. L. Gonçalves, Spherical space forms – homotopy types and self-equivalences for the group $(\mathbb{Z}/a \rtimes \mathbb{Z}/b) \times \mathrm{SL}_2$, *Cand. Math. Bull.* **50** (2) (2007), 206–214.

[GoG5] M. Golasiński and D. L. Gonçalves, Automorphism groups of generalized (binary) Icosahedral, Tetrahedral and octahedral groups, preprint, 2006.

[GoG6] M. Golasiński and D. L. Gonçalves, Spherical space forms – homotopy types and self-equivalences for the group $(\mathbb{Z}/a \rtimes \mathbb{Z}/b) \times \mathrm{TL}_2$, preprint, 2007.

[GG1] D. L. Gonçalves and J. Guaschi, The roots of the full twist for surface braid groups, *Math. Proc. Camb. Phil. Soc.* **137** (2004), 307–320.

[GG2] D. L. Gonçalves and J. Guaschi, The braid group $B_{n,m}(\mathbb{S}^2)$ and the generalised Fadell-Neuwirth short exact sequence, *J. Knot Theory and its Ramifications* **14** (2005), 375–403.

[GG3] D. L. Gonçalves and J. Guaschi, The lower central and derived series of the braid groups $B_n(\mathbb{S}^2)$ and $B_m(\mathbb{S}^2 \setminus \{x_1, \ldots, x_n\})$, preprint, March 2006, `arXiv:math/0603701`.

[GG4] D. L. Gonçalves and J. Guaschi, The quaternion group as a subgroup of the sphere braid groups, *Bull. London Math. Soc.* **39** (2007), 232–234.

[Hm] M.-E. Hamstrom, Homotopy groups of the space of homeomorphisms on a 2-manifold, *Illinois J. Math.* **10** (1966), 563–573.

[Hn] V. L. Hansen, Braids and Coverings: Selected topics, *London Math. Society Student Text* **18**, Cambridge University Press, 1989.

[J] D. L. Johnson, Presentation of groups, LMS Lecture Notes 22 (1976), Cambridge University Press.

[K] S. P. Kerckhoff, The Nielsen realization problem, *Bull. Amer. Math. Soc.* **2** (1980), 452–454.

[Li] V. Lin, Configuration spaces of $\mathbb{C}$ and $\mathbb{CP}^1$: some analytic properties, preprint, March 2004, `arXiv:math.AG/0403120`.

[Ma] W. Magnus, Über Automorphismen von Fundamentalgruppen berandeter Flächen, *Math. Ann.* **109** (1934), 617–646.

[MKS] W. Magnus, A. Karrass and D. Solitar, Combinatorial group theory, reprint of the 1976 second edition, Dover Publications, Inc., Mineola, NY, 2004.

[Mu] K. Murasugi, Seifert fibre spaces and braid groups, *Proc. London Math. Soc.* **44** (1982), 71–84.

[N] M. H. A. Newman, On a string problem of Dirac, *J. London Math. Soc.* **17** (1942), 173–177.

[Sc] G. P. Scott, The space of homeomorphisms of a 2-manifold, *Topology* **9** (1970), 97–109.

[St] M. Stukow, Conjugacy classes of finite subgroups of certain mapping class groups, Seifert fibre spaces and braid groups, *Turkish J. Math.* **2** (2004), 101–110.

[ThC] C. B. Thomas, Elliptic structures on 3-manifolds, Lecture Notes in Mathematics, University of Chicago Department of Mathematics, Chicago, IL, 1983.

[ThJ] J. G. Thompson, Note on $H(4)$, *Comm. Algebra* **22** (1994), 5683–5687.

[VB] J. Van Buskirk, Braid groups of compact 2-manifolds with elements of finite order, *Trans. Amer. Math. Soc.* **122** (1966), 81–97.

[Z] O. Zariski, The topological discriminant group of a Riemann surface of genus p, *Amer. J. Math.* **59** (1937), 335–358.